

ALGEBRAIC AND ANALYTIC DIRAC INDUCTION FOR GRADED AFFINE HECKE ALGEBRAS

DAN CIUBOTARU, ERIC M. OPDAM, AND PETER E. TRAPA

ABSTRACT. We define the algebraic Dirac induction map Ind_D for graded affine Hecke algebras. The map Ind_D is a Hecke algebra analog of the explicit realization of the Baum-Connes assembly map in the K -theory of the reduced C^* -algebra of a real reductive group using Dirac operators. The definition of Ind_D is uniform over the parameter space of the graded affine Hecke algebra. We show that the map Ind_D defines an isometric isomorphism from the space of elliptic characters of the Weyl group (relative to its reflection representation) to the space of elliptic characters of the graded affine Hecke algebra. We also study a related analytically defined global elliptic Dirac operator between unitary representations of the graded affine Hecke algebra which are realized in the spaces of sections of vector bundles associated to certain representations of the pin cover of the Weyl group. In this way we realize all irreducible discrete series modules of the Hecke algebra in the kernels (and indices) of such analytic Dirac operators. This can be viewed as a graded Hecke algebra analogue of the Atiyah-Schmid construction of discrete series for real semisimple Lie groups.

CONTENTS

1. Introduction	1
2. Preliminaries: Euler-Poincaré pairings	5
3. Preliminaries: Dirac operators for the graded affine Hecke algebra	8
4. Dirac induction (algebraic version)	11
5. Orthogonal bases for spaces of virtual elliptic characters	19
6. Dirac induction (analytic version)	27
References	35

1. INTRODUCTION

1.1. Let W be a real reflection group with reflection representation V . Let k be a real valued W -invariant function on the set of simple reflections and let \mathbb{H} be the associated graded affine Hecke algebra with parameters specialized at k (Definition 2.2). In this paper we define the Dirac induction map Ind_D for \mathbb{H} . This map could be thought of as an algebraic analog of (the discrete part of) the explicit realization of the Baum-Connes assembly map $\mu : K_0^G(X) \rightarrow K_0(C_{red}^*(G))$ for the K -theory of the reduced C^* -algebra $C_{red}^*(G)$ of a connected real reductive group G

The authors thank the Max Planck Institute for Mathematics in Bonn for the hospitality and support while part of this research was carried out. In particular, they thank the organizers of “Analysis on Lie groups” for their invitation to participate in the program. This research was supported by nsf-dms 0968065 and NSA-AMS 081022 (D. Ciubotaru), ERC-advanced grant no. 268105 (E. Opdam), nsf-dms 0968275 and 0968060 (P. Trapa).

using Dirac operators (see [La, Section 2.1] for a concise account). In that context $X = G/K$ is the associated Riemannian symmetric space (which we assume to have a G -invariant spin structure) and $K_0^G(X)$ denotes the equivariant K -homology with G -compact supports.

In the present paper we describe the map Ind_D only on the discrete part of the equivariant K -homology with G -compact support of the group $G = V \rtimes W$ acting on $X = V$. After tensoring by \mathbb{C} this space $K_{0, \text{disc}}^G(X)$ can be identified with the complexified space $\overline{R}_{\mathbb{C}}(W)$ of virtual elliptic characters of W (in the sense of Reeder [R], relative to the action of W on V). This space comes equipped with a natural Hermitian inner product, the elliptic pairing. The map Ind_D which we construct is an isometric isomorphism

$$\text{Ind}_D : \overline{R}_{\mathbb{C}}(W) \rightarrow \overline{R}_{\mathbb{C}}(\mathbb{H})$$

where $\overline{R}_{\mathbb{C}}(\mathbb{H})$ is the complexified space of virtual elliptic characters of \mathbb{H} , equipped with its natural Euler-Poincaré pairing. The latter space can be identified canonically with the complexification of the discrete part of the Grothendieck group $K_0(\mathbb{H})$ of finitely generated projective modules over \mathbb{H} , elucidating the analogy with the (discrete part of) the assembly map as realized by Dirac operators. We study the integrality properties of this map, and the central characters of $\text{Ind}_D(\delta)$. An important role in this study is played by the Vogan conjecture as proved in [BCT], and the study of certain irreducible characters of the pin cover of W .

Next we consider a related analytically defined global elliptic Dirac operator between unitary representations of the graded affine Hecke algebra which are realized on the spaces of sections of certain vector bundles associated to a representation of the pin cover of the Weyl group. In this way we realize all irreducible discrete series modules of the Hecke algebra in the kernels (and indices) of such analytic Dirac operators. This can be viewed as a graded Hecke algebra analogue of the Atiyah-Schmid construction of discrete series for real semisimple Lie groups.

The results in this paper provide direct links between three directions of research in the area of affine Hecke algebras:

- (1) the theory of the Dirac operator and Dirac cohomology, as defined for graded affine Hecke algebras in [BCT, C, CT];
- (2) the Euler-Poincaré pairing and elliptic pairing of affine Hecke algebras and Weyl groups, as developed in [OS2, R, So];
- (3) the harmonic analysis approach of [EOS, HO, O1] to the study of unitary modules of graded Hecke algebras, particularly discrete series.

We have used results on the existence and the central support of the discrete series of \mathbb{H} from [O1], [O2], [OS2] in order to define the index of the global elliptic Dirac operators. We hope to replace this by analytic results in the index theory of equivariant elliptic operators in a sequel to the present paper. This could hopefully also shed more light on formal degrees of discrete series representations of \mathbb{H} .

1.2. Let us explain the main results of the paper in more detail. Let $C(V)$ be the Clifford algebra defined with respect to V and a W -invariant inner product $\langle \cdot, \cdot \rangle$ on V . Let \widetilde{W} be the pin double cover of W , a subgroup of the group $\text{Pin}(V)$. When $\dim V$ is odd, $C(V)$ has two nonisomorphic complex simple modules S^+, S^- which remain irreducible when restricted to \widetilde{W} . When $\dim V$ is even, there is a single spin module S of $C(V)$, whose restriction to the even part of $C(V)$ is a sum of

two nonisomorphic simple modules S^+, S^- . The modules S^\pm are irreducible \widetilde{W}' -representations, where \widetilde{W}' is the index two subgroup of \widetilde{W} given by the kernel of the sign representation. In order to describe the results uniformly, denote $\widetilde{W}' = \widetilde{W}$, when $\dim V$ is odd. Let W' be the image of \widetilde{W}' in W under projection.

In [BCT], an analogue of the classical Dirac element is defined: in our case, this is \mathcal{D} , an element of $\mathbb{H} \otimes C(V)$. For every finite dimensional \mathbb{H} -module X , left multiplication by \mathcal{D} gives rise to Dirac operators, which are \widetilde{W}' -invariant:

$$D^\pm : X \otimes S^\pm \rightarrow X \otimes S^\mp. \quad (1.2.1)$$

The Dirac index of X is the virtual \widetilde{W}' -module $I(X) = H_D^+ - H_D^-$, where $H_D^\pm = \ker D^\pm / \ker D^\pm \cap \text{im } D^\mp$ are the Dirac cohomology groups. A standard fact is that $I(X) = X \otimes (S^+ - S^-)$, see Lemma 4.1.

Let $R_{\mathbb{C}}(\mathbb{H})$ be the complex Grothendieck group of \mathbb{H} -modules, and let $\langle \cdot, \cdot \rangle_{\mathbb{H}}^{\text{EP}}$ denote the Euler-Poincaré pairing on $R_{\mathbb{C}}(\mathbb{H})$ (section 2.5). The radical of this form is spanned by parabolically induced modules, and let $\overline{R}_{\mathbb{C}}(\mathbb{H})$ be the quotient by the radical, the space of (virtual) elliptic \mathbb{H} -modules.

The algebra \mathbb{H} contains a copy of the group algebra $\mathbb{C}[W]$ of the Weyl group as a subalgebra. The Grothendieck group $R_{\mathbb{C}}(W)$ has an elliptic pairing $\langle \cdot, \cdot \rangle_W^{\text{ell}}$ defined in [R] whose radical is spanned by induced representations (see section 2.4). Let $\overline{R}_{\mathbb{C}}(W)$ be the quotient by the radical of the form, the space of (virtual) elliptic W -representations. An easy calculation noticed first in [CT], see Theorem 4.2, shows that for every $\delta \in \overline{R}_{\mathbb{Z}}(W)$, there exist associate \widetilde{W}' -representations $\tilde{\delta}^+$ and $\tilde{\delta}^-$ such that $\langle \tilde{\delta}^+, \tilde{\delta}^- \rangle_{\widetilde{W}'} = 0$ and

$$\delta \otimes (S^+ - S^-) = \tilde{\delta}^+ - \tilde{\delta}^-, \text{ and } \langle \delta, \delta \rangle_W^{\text{ell}} = \langle \tilde{\delta}^+, \tilde{\delta}^+ \rangle_{\widetilde{W}'}. \quad (1.2.2)$$

The relation between the elliptic theories of \mathbb{H} and W is given by the restriction map. Precisely, combining results of [OS1, OS2, So] for the affine Hecke algebra and Lusztig's reduction theorems [L1], one sees that the map

$$r : \overline{R}_{\mathbb{C}}(\mathbb{H}) \rightarrow \overline{R}_{\mathbb{C}}(W), \quad [X] \mapsto [X|_W] \quad (1.2.3)$$

is a linear isometry with respect to the pairing $\langle \cdot, \cdot \rangle_{\mathbb{H}}^{\text{EP}}$ and $\langle \cdot, \cdot \rangle_W^{\text{ell}}$. Our first main result is the construction of an inverse Ind_D for the isometry r , which we call the algebraic Dirac induction map. The definition of Ind_D combines elements of K-theory with the Dirac index of \mathbb{H} -modules. Let $K_0(\mathbb{H})_{\mathbb{C}}$ be the complex Grothendieck group of finitely generated projective \mathbb{H} -modules and let $F_0 H_0(\mathbb{H})$ be the subspace of projective modules with zero dimensional support (Definition 4.8). The rank pairing (4.3.1) induces an injective linear map

$$\Phi : F_0 H_0(\mathbb{H}) \rightarrow \overline{R}_{\mathbb{C}}(\mathbb{H}) \quad (1.2.4)$$

with good properties with respect to the rank pairing and the Euler-Poincaré pairing (Lemma 4.9). Then the map Ind_D is defined by extending linearly

$$\text{Ind}_D(\delta) = \Phi([\mathbb{H} \otimes_{W'} ((\delta^+)^* \otimes (S^+ - S^-))]), \quad \delta \in \overline{R}_{\mathbb{Z}}(W). \quad (1.2.5)$$

The following result is part of Theorem 4.11.

Theorem 1.1. *The map Ind_D is well-defined, and it is the inverse of the map r :*

$$\langle \text{Ind}_D(\delta), X \rangle_{\mathbb{H}}^{\text{EP}} = \langle \delta, r(X) \rangle_W^{\text{ell}}, \quad (1.2.6)$$

for every $\delta \in \overline{R}_{\mathbb{C}}(W)$ and $X \in \overline{R}_{\mathbb{C}}(\mathbb{H})$.

As a consequence, we find that whenever δ is a rational multiple of a pure element in $\overline{R}_{\mathbb{Z}}(W)$, see Definition 4.12, then $\text{Ind}_D(\delta)$ is supported in a single central character $\Lambda(\delta)$, see Corollaries 4.13 and 5.7. We also prove in Theorem 4.18, that the central character $\text{Ind}_D(\delta)$ depends linearly in the parameter function k of the Hecke algebra.

In section 5, we study further the map Ind_D in the case when the root system R is irreducible, in particular, we investigate its behaviour with respect to the integral lattices $\overline{R}_{\mathbb{Z}}(W)$ and $\overline{R}_{\mathbb{Z}}(\mathbb{H})$, see section 5.5. To this end, in Theorem 5.1 we find appropriate orthogonal bases consisting of pure elements for the lattices $\overline{R}_{\mathbb{Z}}(\mathbb{H})$ and $\overline{R}_{\mathbb{Z}}(W)$.

1.3. The second part of the paper concerns a realization of discrete series \mathbb{H} -modules in the index (hence kernel) of certain global Dirac operators. This can be regarded as a Hecke algebra analogue of the construction in [AS]. One complication in our setting is that, while the discrete series and tempered spectra of the graded affine Hecke algebra \mathbb{H} are known [L2, O1], there is no known abstract Plancherel formula for \mathbb{H} . To bypass this difficulty we construct directly certain analytic models $\mathcal{X}'_{\omega}(E \otimes S^{\pm})$, which are (pre)unitary left \mathbb{H} -modules, for every \widetilde{W}' -representation E . It is easy to check that every irreducible \mathbb{H} -submodule of $\mathcal{X}'_{\omega}(E \otimes S^{\pm})$ is necessarily a discrete series (Lemma 6.14).

These models are analogous with the spaces of sections of spinor bundles for Riemannian symmetric spaces from [AS]. The construction that we use is an adaptation of the ones from [HO] and [EOS] (the latter being in the setting of the trigonometric Cherednik algebra). A new ingredient is an \mathbb{H} -invariant inner product, this is Theorem 6.3, a generalization of the unitary structure from [HO]. Once these definitions are in place, we can consider global Dirac operators acting “on the right”:

$$D_E^{\pm} : \mathcal{X}'_{\omega}(E \otimes S^{\pm}) \rightarrow \mathcal{X}'_{\omega}(E \otimes S^{\mp}). \quad (1.3.1)$$

If λ is a central character of \mathbb{H} , restrict D_E^{\pm} to the subspace $\mathcal{X}'_{\omega}(E \otimes S^{\pm})_{\lambda}$ of $\mathcal{X}'_{\omega}(E \otimes S^{\pm})$ on which the center $Z(\mathbb{H})$ acts via λ . Denote by $D_E^{\pm}(\lambda)$ the restricted Dirac operators. Define the global Dirac index to be the formal expression

$$I_E = \bigoplus_{\lambda} (\ker D_E^{+}(\lambda) - \ker D_E^{-}(\lambda)), \quad (1.3.2)$$

a virtual left \mathbb{H} -module. We show in section 6.5 that this sum is finite.

We define an analytic Dirac induction map $\text{Ind}_D^{\omega} : \overline{R}_{\mathbb{C}}(W) \rightarrow \overline{R}_{\mathbb{C}}(\mathbb{H})$ as follows. Let $\delta \in \overline{R}_{\mathbb{Z}}(W)$ be given, and let δ^{+} be the \widetilde{W}' -representation in (1.2.2). Set

$$\text{Ind}_D^{\omega}(\delta) = I_{(\delta^{+})^{*}}. \quad (1.3.3)$$

A consequence of Vogan’s conjecture (Theorem 3.2) is that the irreducible \mathbb{H} -modules that occur in $\text{Ind}_D^{\omega}(\delta)$ must have a prescribed central character $\Lambda(\delta)$, see Theorem 4.18. The main results about Ind_D^{ω} can be summarized as follows (see Theorem 6.16).

Theorem 1.2. (1) *For every irreducible \mathbb{H} -module X , $\text{Hom}_{\mathbb{H}}(X, \text{Ind}_D^{\omega}(\delta)) = 0$ unless X is a discrete series with central character $\Lambda(\delta)$, in which case*

$$\dim \text{Hom}_{\mathbb{H}}(X, \text{Ind}_D^{\omega}(\delta)) = \langle \mathbf{r}(X), \delta \rangle_W^{\text{ell}}. \quad (1.3.4)$$

- (2) Let X be an irreducible discrete series \mathbb{H} -module. Then $X \cong \text{Ind}_D^\omega(r(X))$ as \mathbb{H} -modules. In particular, the image of the map Ind_D^ω is the span of the discrete spectrum of \mathbb{H} .

Notation 1.3. If k is a unital commutative ring and A is a k -associative algebra, denote by $\text{Rep}(A)$ the category of (left) A -modules and by $\text{Rep}_{\text{fd}}(A)$ the category of finite dimensional modules; if G is a group, denote by $\text{Rep}_k(G)$, the category of $k[G]$ -modules. Let $R_K(A)$ and $R_K(G)$ denote the Grothendieck rings of $\text{Rep}(A)$ and $\text{Rep}_k(G)$, respectively, with coefficients in a ring K .

2. PRELIMINARIES: EULER-POINCARÉ PAIRINGS

2.1. The root system. Fix a semisimple real root system $\Phi = (V, R, V^\vee, R^\vee)$. In particular, V and V^\vee are finite dimensional real vector spaces, $R \subset V \setminus \{0\}$ generates V , $R^\vee \subset V^\vee \setminus \{0\}$ generates V^\vee and there is a perfect bilinear pairing

$$(\cdot, \cdot) : V \times V^\vee \rightarrow \mathbb{R},$$

which induces a bijection between R and R^\vee such that $(\alpha, \alpha^\vee) = 2$ for all $\alpha \in R$. For every $\alpha \in R$, let $s_\alpha : V \rightarrow V$ denote the reflection about the root α given by $s_\alpha(v) = v - (v, \alpha^\vee)\alpha$ for all $v \in V$. We also identify s_α with the map $s_\alpha : V^\vee \rightarrow V^\vee$, $s_\alpha(v') = v' - (\alpha, v')\alpha^\vee$.

Let W be the subgroup of $GL(V)$ generated by $\{s_\alpha : \alpha \in R\}$; we may also regard W as a subgroup of $GL(V^\vee)$. Fix a basis F of R and set $S = \{s_\alpha : \alpha \in F\}$. Then (W, S) is a finite Coxeter group.

Let $Q = \mathbb{Z}R \subset V$ denote the root lattice, $\mathcal{P} \subset V$ the weight lattice, and form the affine Weyl group $W_{\text{aff}} = Q \rtimes W$. Let F_{aff} be the set of simple affine roots, and $S_{\text{aff}} \supset S$ the corresponding set of simple affine reflections. Let ℓ denote the length function of the Coxeter group $(W_{\text{aff}}, S_{\text{aff}})$.

The complexified vector spaces will be denoted by $V_{\mathbb{C}}$ and $V_{\mathbb{C}}^\vee$.

2.2. The affine Hecke algebra \mathcal{H} . Let $q : S_{\text{aff}} \rightarrow \mathbb{R}_{>0}$ be a function such that $q_s = q_{s'}$ whenever $s, s' \in S_{\text{aff}}$ are W_{aff} -conjugate. For every $s \in S_{\text{aff}}$, let $q_s^{1/2}$ denote the positive square root of q_s .

Definition 2.1. The affine Hecke algebra $\mathcal{H} = \mathcal{H}(W_{\text{aff}}, q)$ is the unique associative unital free \mathbb{C} -algebra with basis $\{T_w : w \in W_{\text{aff}}\}$ subject to the relations:

- (i) $T_w T_{w'} = T_{ww'}$ if $\ell(ww') = \ell(w) + \ell(w')$;
- (ii) $(T_s - q_s^{1/2})(T_s + q_s^{-1/2}) = 0$, for all $s \in S_{\text{aff}}$.

A particular instance is when the parameters q_s of \mathcal{H} are specialized to 1; then \mathcal{H} becomes $\mathbb{C}[W_{\text{aff}}]$.

A result of Bernstein (*cf.* [L1, Proposition 3.11]) says that the center $Z(\mathcal{H})$ of \mathcal{H} is isomorphic with the algebra of W -invariant complex functions on Q . In particular, the central characters, i.e., the homomorphisms $\chi^t : Z(\mathcal{H}) \rightarrow \mathbb{C}$ are parameterized by classes Wt in $W \backslash T$, where T is the complex torus $\mathbb{C} \otimes_{\mathbb{Z}} \mathcal{P}$. We say that the central character χ^t is real if $t \in \mathbb{R} \otimes_{\mathbb{Z}} \mathcal{P}$.

Denote by $\text{Rep}(\mathcal{H})_0$ and $\text{Rep}_{\text{fd}}(\mathcal{H})_0$ the respective full subcategories of \mathcal{H} -modules with real central characters. If $P \subset F$, let W_P denote the subgroup of W generated by $\{s_\alpha : \alpha \in P\}$. Let \mathcal{H}_P be the parabolic affine Hecke algebra, see [So, section 1.4] for example.

Delorme-Opdam defined the Schwartz algebra \mathcal{S} of \mathcal{H} ([DO, section 2.8]) by

$$\mathcal{S} = \left\{ \sum_{w \in W_{\text{aff}}} c_w T_w : \text{for all } n \in \mathbb{N}, (1 + \ell(w))^n c_w \text{ is bounded as a function in } w \in W_{\text{aff}} \right\}. \quad (2.2.1)$$

Then \mathcal{S} is a nuclear Fréchet algebra (in the sense of [O1, Definition 6.6]) with respect to the family of norms

$$p_n \left(\sum_{w \in W_{\text{aff}}} c_w T_w \right) := \sup \{ (1 + \ell(w))^n |c_w| : w \in W_{\text{aff}} \}.$$

An irreducible module $X \in \text{Rep}_{\mathbb{C}}(\mathcal{H})$ is called tempered if it can be extended to a \mathcal{S} -module. The irreducible summands in the \mathcal{H} -decomposition of \mathcal{S} are called discrete series. These definitions of tempered and discrete series agree with those given by the Casselman criterion which we do not recall here, instead we refer to [O1, section 2.7] for the details. Let $\text{Rep}(\mathcal{S})$ denote the category of \mathcal{S} -representations.

2.3. The graded affine Hecke algebra \mathbb{H} . Let $\underline{k} = \{k_{\alpha} : \alpha \in F\}$ be a set of indeterminates such that $k_{\alpha} = k_{\alpha'}$ whenever α, α' are W -conjugate. Denote $A = \mathbb{C}[\underline{k}]$. Let $\mathbb{C}[W]$ denote the group algebra of W and $S(V_{\mathbb{C}})$ the symmetric algebra over $V_{\mathbb{C}}$. The group W acts on $S(V_{\mathbb{C}})$ by extending the action on V . For every $\alpha \in F$, denote the difference operator by

$$\Delta : S(V_{\mathbb{C}}) \rightarrow S(V_{\mathbb{C}}), \quad \Delta_{\alpha}(p) = \frac{p - s_{\alpha}(p)}{\alpha}, \text{ for all } p \in S(V_{\mathbb{C}}). \quad (2.3.1)$$

Definition 2.2. The generic graded affine Hecke algebra $\mathbb{H}_A = \mathbb{H}(\Phi, F, \underline{k})$ is the unique associative unital A -algebra such that

- (i) $\mathbb{H}_A = S(V_{\mathbb{C}}) \otimes A[W]$ as a $(S(V_{\mathbb{C}}), A[W])$ -bimodule;
- (ii) $s_{\alpha} \cdot p = s_{\alpha}(p) \cdot s_{\alpha} + k_{\alpha} \Delta_{\alpha}(p)$, for all $\alpha \in F$, $p \in S(V_{\mathbb{C}})$.

If $k : F \rightarrow \mathbb{R}_{\geq 0}$ is a function such that $k_{\alpha} = k_{\alpha'}$ whenever $\alpha, \alpha' \in F$ are W -conjugate, let \mathbb{H} (or \mathbb{H}_k , when we wish to emphasize the dependence on the parameter function k) be the specialization of \mathbb{H}_A at k , i.e., $\mathbb{H} = \mathbb{C}_k \otimes_A \mathbb{H}_A$, where \mathbb{C}_k is the A -module on which \underline{k} acts by k .

A result of Lusztig [L1, Proposition 4.5] says that the center $Z(\mathbb{H}_A)$ of \mathbb{H}_A is $A \otimes_{\mathbb{C}} S(V_{\mathbb{C}})^W$. In particular, $Z(\mathbb{H}) = S(V_{\mathbb{C}})^W$ and the central characters $\chi^{\lambda} : Z(\mathbb{H}) \rightarrow \mathbb{C}$ are parameterized by classes $W\lambda$ in $W \backslash V_{\mathbb{C}}^{\vee}$. We say that the central character χ^{λ} is real if $\lambda \in V^{\vee}$.

Denote by $\text{Rep}(\mathbb{H})_0$ and $\text{Rep}_{\text{fd}}(\mathbb{H})_0$ the respective full subcategories of \mathbb{H} -modules with real central characters. If $P \subset F$, let \mathbb{H}_P be the parabolic graded affine Hecke subalgebra, see [So, section 1.4] for example.

Definition 2.3. An \mathbb{H} -module X is called tempered if every $S(V_{\mathbb{C}})$ -weight $\nu \in V_{\mathbb{C}}^{\vee}$ of X satisfies the Casselman criterion

$$(\omega, \Re \nu) \leq 0, \text{ for all fundamental weights } \omega \in \mathcal{P}. \quad (2.3.2)$$

The module X is called a discrete series if all the inequalities in (2.3.2) are strict. Denote by $\text{DS}(\mathbb{H})$ the set of irreducible discrete series \mathbb{H} -modules.

A particular case of Lusztig's reduction theorems is the following.

Theorem 2.4. *There is an equivalence of categories*

$$\eta : \text{Rep}_{\text{fd}}(\mathcal{H}(W_{\text{aff}}, q))_0 \rightarrow \text{Rep}_{\text{fd}}(\mathbb{H}_k)_0,$$

where the relation between the parameters q and k is as in [OS2, (26)]. Moreover, $X \in \text{Rep}_{\text{fd}}(\mathcal{H}(W_{\text{aff}}, q))_0$ is irreducible tempered (resp. discrete series) if and only if $\eta(X) \in \text{Rep}_{\text{fd}}(\mathbb{H}_k)_0$ is irreducible tempered (resp. discrete series).

2.4. Euler-Poincaré pairing for the Weyl group. If $\sigma \in R_{\mathbb{C}}(W)$, let χ_{σ} denote the character of σ .

Definition 2.5 ([R, section 2.1]). For $\sigma, \mu \in R_{\mathbb{C}}(W)$, the Euler-Poincaré pairing is the Hermitian form

$$\langle \sigma, \mu \rangle_W^{\text{ell}} = \frac{1}{|W|} \sum_{w \in W} \overline{\chi_{\sigma}(w)} \chi_{\mu}(w) \det_{V_{\mathbb{C}}}(1 - w).$$

An element $w \in W$ is called elliptic if $\det_{V_{\mathbb{C}}}(1 - w) \neq 0$.

For every parabolic subgroup W_P of W corresponding to $P \subset F$, let $\text{ind}_{W_P}^W : \text{Rep}_{\mathbb{C}}(W_P) \rightarrow \text{Rep}_{\mathbb{C}}(W)$ denote the induction functor.

Theorem 2.6 ([R, (2.1.1), (2.2.2)]). *The radical of the form $\langle \cdot, \cdot \rangle_W^{\text{ell}}$ on $R_{\mathbb{C}}(W)$ is $\sum_{P \subsetneq F} \text{ind}_{W_P}^W(R_{\mathbb{C}}(W_P))$. Moreover:*

- (i) $\langle \cdot, \cdot \rangle_W^{\text{ell}}$ induces a positive definite Hermitian form on

$$\overline{R}_{\mathbb{C}}(W) = R_{\mathbb{C}}(W) / \sum_{P \subsetneq F} \text{ind}_{W_P}^W(R_{\mathbb{C}}(W_P)).$$

- (ii) *the dimension of $\overline{R}_{\mathbb{C}}(W)$ equals the number of elliptic conjugacy classes in W .*

2.5. Euler-Poincaré pairing for \mathcal{H} .

Definition 2.7 ([OS1, (3.15), Theorem 3.5]). Define on $R_{\mathbb{C}}(\mathcal{H})$ a Hermitian form $\langle \cdot, \cdot \rangle_{\mathcal{H}}^{\text{EP}}$ by

$$\langle X, Y \rangle_{\mathcal{H}}^{\text{EP}} = \sum_{i \geq 0} (-1)^i \dim \text{Ext}_{\mathcal{H}}^i(X, Y), \text{ for all } X, Y \in R_{\mathbb{C}}(\mathcal{H}).$$

Let $\text{ind}_{\mathcal{H}_P}^{\mathcal{H}} : \text{Rep}_{\mathbb{C}}(\mathcal{H}_P) \rightarrow \text{Rep}_{\mathbb{C}}(\mathcal{H})$ denote the induction functor. If $P \subsetneq F$, one can see that the space $\text{ind}_{\mathcal{H}_P}^{\mathcal{H}}(R_{\mathbb{C}}(\mathcal{H}_P))$ is in the radical of $\langle \cdot, \cdot \rangle_{\mathcal{H}}^{\text{EP}}$, [OS1, Proposition 3.4] Therefore, $\langle \cdot, \cdot \rangle_{\mathcal{H}}^{\text{EP}}$ factors through

$$\overline{R}_{\mathbb{C}}(\mathcal{H}) = R_{\mathbb{C}}(\mathcal{H}) / \sum_{P \subsetneq F} \text{ind}_{\mathcal{H}_P}^{\mathcal{H}}(R_{\mathbb{C}}(\mathcal{H}_P)).$$

One can define the pairing $\langle \cdot, \cdot \rangle_{\mathcal{S}}^{\text{EP}}$ in $R_{\mathbb{C}}(\mathcal{S})$ and consider the space of (virtual) elliptic tempered modules $\overline{R}_{\mathbb{C}}(\mathcal{S})$. As a consequence of the Langlands classification, one sees easily that $\overline{R}_{\mathbb{C}}(\mathcal{H}) = \overline{R}_{\mathbb{C}}(\mathcal{S})$. The fact that this isomorphism is an isometry with respect to the Euler-Poincaré pairings follows from [OS1, Corollary 3.7]:

$$\text{Ext}_{\mathcal{H}}^i(X, Y) \cong \text{Ext}_{\mathcal{S}}^i(X, Y), \text{ for all finite dimensional tempered } \mathcal{H}\text{-modules } X, Y. \quad (2.5.1)$$

Theorem 2.8 ([OS1, Theorem 3.8]). *If X is an irreducible discrete series \mathcal{H} -module and Y is a finite dimensional tempered \mathcal{H} -module, then*

$$\mathrm{Ext}_{\mathcal{H}}^i(X, Y) = \mathrm{Ext}_{\mathcal{S}}^i(X, Y) = 0, \text{ for all } i > 0. \quad (2.5.2)$$

In particular, $\langle X, Y \rangle_{\mathcal{H}}^{\mathrm{EP}} = 1$ if $X \cong Y$ and 0 otherwise.

Solleveld [So, Theorem 4.4.2] defines a scaling map

$$\tau_0 : R_{\mathbb{C}}(\mathcal{S}) \rightarrow R_{\mathbb{C}}(W_{\mathrm{aff}}), \quad \tau_0(X) = \lim_{\epsilon \rightarrow 0} X|_{q_s \rightarrow q_s^\epsilon}; \quad (2.5.3)$$

this has the property that it factors through $\overline{R}_{\mathbb{C}}(\mathcal{S}) = \overline{R}_{\mathbb{C}}(\mathcal{H}) \rightarrow \overline{R}_{\mathbb{C}}(W_{\mathrm{aff}})$. The relation with the elliptic theory of the Weyl group can be summarized in the following statements which are a combination of [OS1, Theorem 3.2, Proposition 3.9] together with [OS2, Corollary 7.4] and [So, Theorem 2.3.1].

- Theorem 2.9** (Opdam-Solleveld). (i) *The map τ_0 defines a linear isomorphism $\overline{R}_{\mathbb{C}}(\mathcal{H}) \rightarrow \overline{R}_{\mathbb{C}}(W_{\mathrm{aff}})$, $[X] \mapsto [\tau_0(X)]$.*
(ii) *The isomorphism τ_0 from (i) is an isometry with respect to $\langle \cdot, \cdot \rangle_{\mathcal{H}}^{\mathrm{EP}}$ and $\langle \cdot, \cdot \rangle_{W_{\mathrm{aff}}}^{\mathrm{EP}}$, respectively.*
(iii) *The map $\tau_{\mathrm{ell}} : \overline{R}_{\mathbb{C}}(\mathcal{H})_0 \rightarrow \overline{R}_{\mathbb{C}}(W)$, given by $[X] \mapsto [\tau_0(X)|_W]$ is a linear isomorphism and an isometry with respect to $\langle \cdot, \cdot \rangle_{\mathcal{H}}^{\mathrm{EP}}$ and $\langle \cdot, \cdot \rangle_W^{\mathrm{ell}}$.*

2.6. Euler-Poincaré pairing for \mathbb{H} . We need to translate the previous results in the setting of the graded Hecke algebra \mathbb{H} . Define the pairing $\langle \cdot, \cdot \rangle_{\mathbb{H}}^{\mathrm{EP}}$ on $R_{\mathbb{C}}(\mathbb{H})$ and the space of virtual elliptic modules $\overline{R}_{\mathbb{C}}(\mathbb{H})$ in the same way as for \mathcal{H} . Lusztig's reduction equivalence $\eta : \mathrm{Rep}_{\mathrm{fd}}(\mathcal{H})_0 \rightarrow \mathrm{Rep}_{\mathrm{fd}}(\mathbb{H})_0$ induces a linear isomorphism $\overline{\eta} : \overline{R}_{\mathbb{C}}(\mathcal{H})_0 \rightarrow \overline{R}_{\mathbb{C}}(\mathbb{H})_0$ which is an isometry with respect to the Euler-Poincaré pairings. In fact, $\overline{R}_{\mathbb{C}}(\mathbb{H})_0 = \overline{R}_{\mathbb{C}}(\mathbb{H})$, i.e., all finite dimensional elliptic \mathbb{H} -modules have real central character. Comparing with the results recalled before, we arrive at the following corollaries.

Corollary 2.10. *If X is an irreducible discrete series \mathbb{H} -module, and Y is a finite dimensional tempered \mathbb{H} -module, then $\langle X, Y \rangle_{\mathbb{H}}^{\mathrm{EP}} = 1$ if $X \cong Y$ and 0 otherwise.*

Corollary 2.11. *The restriction map $\mathrm{res} : \mathrm{Rep}_{\mathbb{C}}(\mathbb{H}) \rightarrow \mathrm{Rep}_{\mathbb{C}}(W)$, $\mathrm{res}(X) = X|_W$ induces a linear isometry*

$$r : \overline{R}_{\mathbb{C}}(\mathbb{H}) \rightarrow \overline{R}_{\mathbb{C}}(W) \quad (2.6.1)$$

with respect to $\langle \cdot, \cdot \rangle_{\mathbb{H}}^{\mathrm{EP}}$ and $\langle \cdot, \cdot \rangle_W^{\mathrm{ell}}$, respectively. In particular, if X is an irreducible discrete series \mathbb{H} -module, and Y is a finite dimensional tempered \mathbb{H} -module, $\langle r(X), r(Y) \rangle_W^{\mathrm{ell}} = 1$ if $X \cong Y$ and 0 otherwise.

3. PRELIMINARIES: DIRAC OPERATORS FOR THE GRADED AFFINE HECKE ALGEBRA

We retain the notation from the previous section. We recall the construction and basic facts about the Dirac operator for \mathbb{H} .

3.1. The pin cover of the Weyl group. Fix a W -invariant inner product $\langle \cdot, \cdot \rangle$ on V and let $C(V)$ denote the Clifford algebra, the quotient of the tensor algebra of V by the ideal generated by $\{\omega \otimes \omega' + \omega' \otimes \omega + 2\langle \omega, \omega' \rangle : \omega, \omega' \in V\}$. Let $O(V)$ denote the group of orthogonal transformations of V with respect to $\langle \cdot, \cdot \rangle$. We have $W \subset O(V)$. The action of $-1 \in O(V)$ on $C(V)$ induces a $\mathbb{Z}/2\mathbb{Z}$ -grading $C(V) = C(V)_{\mathrm{even}} + C(V)_{\mathrm{odd}}$, and let ϵ be the automorphism of $C(V)$ which is 1

on $C(V)_{\text{even}}$ and -1 on $C(V)_{\text{odd}}$. Let t be the transpose automorphism of $C(V)$ defined by $\omega^t = -\omega$, $\omega \in V$, and $(ab)^t = b^t a^t$ for $a, b \in C(V)$. The pin group is

$$\text{Pin}(V) = \{a \in C(V) : \epsilon(a)Va^{-1} \subset V, a^t = a^{-1}\}; \quad (3.1.1)$$

it is a central $\mathbb{Z}/2\mathbb{Z}$ -extension of $O(V)$:

$$1 \rightarrow \{\pm 1\} \rightarrow \text{Pin}(V) \xrightarrow{p} O(V) \rightarrow 1,$$

where p is the projection $p(a)(\omega) = \epsilon(a)\omega a^{-1}$. Construct the central $\mathbb{Z}/2\mathbb{Z}$ -extension $\widetilde{W} = p^{-1}(W)$ of W :

$$1 \rightarrow \{\pm 1\} \rightarrow \widetilde{W} \xrightarrow{p} W \rightarrow 1. \quad (3.1.2)$$

The group \widetilde{W} has a Coxeter presentation similar to that of W , see [M]:

$$\widetilde{W} = \langle z, \widetilde{s}_\alpha, \alpha \in F : z^2 = 1, \widetilde{s}_\alpha^2 = z, (\widetilde{s}_\alpha \widetilde{s}_\beta)^{m(\alpha, \beta)} = 1, \alpha \neq \beta \in F \rangle. \quad (3.1.3)$$

With this presentation, the embedding of \widetilde{W} in $\text{Pin}(V)$ is given by:

$$z \mapsto -1, \quad \widetilde{s}_\alpha \mapsto \frac{1}{|\alpha|} \alpha. \quad (3.1.4)$$

3.2. The Dirac element. The generic Hecke algebra \mathbb{H}_A (Definition 2.2) has a natural $*$ -operation coming from the relation with the affine Hecke algebra \mathcal{H} and p -adic groups. On generators, this is defined by

$$\begin{aligned} \underline{k}_\alpha^* &= \underline{k}_\alpha, \quad \alpha \in F; \quad w^* = w^{-1}, \quad w \in W; \\ \xi^* &= -w_0 \cdot w_0(\xi) \cdot w_0 = -\xi + \sum_{\beta \in R^+} \underline{k}_\beta(\xi, \beta^\vee) s_\beta, \quad \xi \in V. \end{aligned} \quad (3.2.1)$$

For every $\xi \in V$, define

$$\widetilde{\xi} = \xi - T_\xi, \quad \text{where } T_\xi = \frac{1}{2} \sum_{\beta \in R^+} \underline{k}_\beta(\xi, \beta^\vee) s_\beta \in \mathbb{H}_A. \quad (3.2.2)$$

Then $\widetilde{\xi}^* = -\widetilde{\xi}$, for all $\xi \in V$.

Definition 3.1. Let $\{\xi_i\}, \{\xi^i\}$ be dual bases of $V, \langle \cdot, \cdot \rangle$. The Dirac element is

$$\mathcal{D} = \sum_i \widetilde{\xi}_i \otimes \xi^i \in \mathbb{H}_A \otimes C(V).$$

It does not depend on the choice of bases.

Write ρ for the diagonal embedding of $\mathbb{C}[\widetilde{W}]$ into $\mathbb{H}_A \otimes C(V)$ defined by extending linearly $\rho(\widetilde{w}) = p(\widetilde{w}) \otimes \widetilde{w}$. By [BCT, Lemma 3.4], we see that \mathcal{D} is $\text{sgn } \widetilde{W}$ -invariant, i.e.,

$$\rho(\widetilde{w})\mathcal{D} = \text{sgn}(\widetilde{w})\mathcal{D}\rho(\widetilde{w}), \quad \text{for all } \widetilde{w} \in \widetilde{W}. \quad (3.2.3)$$

Moreover, [BCT, Theorem 3.5] computes \mathcal{D}^2 , and in particular shows that \mathcal{D}^2 acts diagonally on \widetilde{W} -isotypic components of $X \otimes S$, for every irreducible \mathbb{H} -module X and every $C(V)$ -module S . More precisely, define

$$\Omega = \sum_i \xi_i \xi^i \in Z(\mathbb{H}_A), \quad (3.2.4)$$

and

$$\Omega_{\widetilde{W}} = \frac{z}{4} \sum_{\alpha > 0, \beta > 0, s_\alpha(\beta) < 0} \underline{k}_\alpha \underline{k}_\beta |\alpha^\vee| |\beta^\vee| \widetilde{s}_\alpha \widetilde{s}_\beta \in \mathbb{C}[\widetilde{W}]^{\widetilde{W}}. \quad (3.2.5)$$

Then, we have

$$\mathcal{D}^2 = -\Omega \otimes 1 + \rho(\Omega_{\widetilde{W}}). \quad (3.2.6)$$

3.3. Vogan's conjecture. In this section, we sharpen the statement of Vogan's conjecture (proved in [BCT]) related to the Dirac cohomology of \mathbb{H} -modules. The following result is the generic analogue of [BCT, Theorem 4.2].

Theorem 3.2. *Let \mathbb{H}_A be the generic graded Hecke algebra over $A = \mathbb{C}[\underline{k}]$ (see Definition 2.2). Let $z \in Z(\mathbb{H}_A)$ be given. There exists $a \in \mathbb{H}_A \otimes C(V)_{\text{odd}}$ and a unique element $\zeta(z)$ in the center of $A[\widetilde{W}]$ such that*

$$z \otimes 1 = \rho(\zeta(z)) + \mathcal{D}a + a\mathcal{D}, \quad (3.3.1)$$

as elements in $\mathbb{H}_A \otimes C(V)$. Moreover, the map $z \rightarrow \zeta(z)$ defines an algebra homomorphism $\zeta : Z(\mathbb{H}_A) \rightarrow A[\widetilde{W}]^{\widetilde{W}}$.

Notice that under the homomorphism ζ we have

$$\zeta(\Omega) = \Omega_{\widetilde{W}}, \quad (3.3.2)$$

where Ω is as in (3.2.4) and $\Omega_{\widetilde{W}}$ is as in (3.2.5).

Proof. We proceed as in [BCT, section 4]. Let $\mathbb{H}_A^0 \subset \mathbb{H}_A^1 \subset \dots \mathbb{H}_A^n \subset \dots$ be the filtration coming from the degree filtration of $S(V_{\mathbb{C}})$. (The group algebra $A[W]$ has degree zero.) The associated graded object $\bigoplus_j \overline{\mathbb{H}}_A^j$, $\overline{\mathbb{H}}_A^j = \mathbb{H}_A^j / \mathbb{H}_A^{j-1}$, is naturally isomorphic as an A -algebra to $\mathbb{H}_{A,0} = A \otimes_{\mathbb{C}} \mathbb{H}_0$. Define

$$d : \mathbb{H}_A \otimes C(V) \rightarrow \mathbb{H}_A \otimes C(V),$$

by extending linearly $d(h \otimes c_1 \dots c_\ell) = \mathcal{D} \cdot (h \otimes c_1 \dots c_\ell) - (-1)^\ell (h \otimes c_1 \dots c_\ell) \cdot \mathcal{D}$, $h \in \mathbb{H}_A$, $c_i \in V$, and restrict d to

$$d^{\text{triv}} : (\mathbb{H}_A \otimes C(V))^{\text{triv}} \rightarrow (\mathbb{H}_A \otimes C(V))^{\text{sgn}},$$

as in [BCT, section 5]. The statement of Theorem 3.2 follows from

$$\ker(d^{\text{triv}}) = \text{im}(d^{\text{sgn}}) \oplus \rho(A[\widetilde{W}]^{\widetilde{W}}),$$

see [BCT, Theorem 5.1]. This in turn is proved as follows. Firstly, one verifies that $\ker(d^{\text{triv}}) \supset \text{im}(d^{\text{sgn}}) \oplus \rho(A[\widetilde{W}]^{\widetilde{W}})$. Secondly, d^{triv} induces a graded differential

$$\overline{d}^{\text{triv}} : (\mathbb{H}_{A,0} \otimes C(V))^{\text{triv}} \rightarrow (\mathbb{H}_{A,0} \otimes C(V))^{\text{sgn}},$$

for which [BCT, Corollary 5.9] shows that

$$\ker(\overline{d}^{\text{triv}}) = \text{im}(\overline{d}^{\text{sgn}}) \oplus \overline{\rho}(A[\widetilde{W}]^{\widetilde{W}}). \quad (3.3.3)$$

Thirdly, one proceeds by induction on degree in $\mathbb{H}_A \otimes C(V)$ to deduce from (3.3.3) the opposite inclusion $\ker(d^{\text{triv}}) \subset \text{im}(d^{\text{sgn}}) \oplus \rho(A[\widetilde{W}]^{\widetilde{W}})$. This is the step that we need to examine more closely.

Let $b \in \ker(d^{\text{triv}})$ be an element of $\mathbb{H}_A^n \otimes C(V)$ (i.e., an element of degree n in the filtration), which can be specialized to $z \otimes 1$. Since $d^{\text{triv}}(b) = 0$, taking the graded objects, we have $\overline{d}^{\text{triv}}(\overline{b}) = 0$ in $\mathbb{H}_{A,0} \otimes C(V)$. From (3.3.3), there exists $\overline{c} \in (\mathbb{H}_{A,0}^{n-1} \otimes C(V))^{\text{sgn}}$ and $s \in A[\widetilde{W}]^{\widetilde{W}}$ such that

$$\overline{b} = \overline{d}^{\text{sgn}} \overline{c} + \overline{\rho}(s).$$

Choose $c \in (\mathbb{H}_A^{n-1} \otimes C(V))^{\text{sgn}}$ such that \bar{c} is the image of c in $(\mathbb{H}_{A,0}^{n-1} \otimes C(V))^{\text{sgn}}$. For example, if $\bar{c} = \sum w \xi_{w,1} \dots \xi_{w,n-1} \otimes f_w$, $w \in W$, $\xi_{w,i} \in V_{\mathbb{C}}$, $f_w \in C(V)$, we can choose $c = \sum w \tilde{\xi}_{w,1} \dots \tilde{\xi}_{w,n-1} \otimes f_w$. Then

$$\overline{b - d^{\text{sgn}} c - \rho(s)} = \bar{b} - \bar{d}^{\text{sgn}} \bar{c} - \bar{\rho}(s) = 0,$$

hence $b - d^{\text{sgn}} c - \rho(s) \in (\mathbb{H}_A^{n-1} \otimes C(V))^{\text{triv}}$. On the other hand,

$$d^{\text{triv}}(b - d^{\text{sgn}} c - \rho(s)) = d^{\text{triv}}(b) - d^2(c) - d^{\text{triv}}(\rho(s)) = 0,$$

where $d^{\text{triv}}(b) = 0$ by assumption, $d^2(c) = 0$ by [BCT, Lemma 5.3], and $d^{\text{triv}}(\rho(s)) = 0$ by [BCT, Lemma 5.2]. But then, by induction, $b - d^{\text{sgn}} c - \rho(s) = d^{\text{sgn}} c' + \rho(s')$, where $s' \in A[\widetilde{W}]^{\widetilde{W}}$ and $c' \in (\mathbb{H}_A \otimes C(V))^{\text{sgn}}$. \square

Corollary 3.3. *Let $\zeta : Z(\mathbb{H}_A) \rightarrow A[\widetilde{W}]^{\widetilde{W}}$ be the algebra homomorphism from Theorem 3.2. Then the image of ζ lies in $A[\ker \text{sgn}]$, where sgn is the sign \widetilde{W} -representation.*

Proof. For every $z \in Z(\mathbb{H}_A)$, consider the defining equation (3.3.1) for $\zeta(z)$. Since $a \in \mathbb{H}_A \otimes C(V)_{\text{odd}}$, we have $\mathcal{D}a + a\mathcal{D} \in \mathbb{H}_A \otimes C(V)_{\text{even}}$, and also clearly $z \otimes 1 \in \mathbb{H}_A \otimes C(V)_{\text{even}}$. Thus $\rho(\zeta(z)) \in \mathbb{H}_A \otimes C(V)_{\text{even}}$, and the claim follows. \square

4. DIRAC INDUCTION (ALGEBRAIC VERSION)

The goal of this section is to construct an inverse of the restriction map r from (2.6.1), using the Dirac index theory.

4.1. The local Dirac index. The notion of Dirac index for finite dimensional \mathbb{H} -modules was introduced in [CT, section 2.9]. Let X be a finite dimensional \mathbb{H} -module. Denote

$$\widetilde{W}' = \begin{cases} \widetilde{W}, & \text{if } \dim(V) \text{ is odd,} \\ \ker \text{sgn}, & \text{if } \dim(V) \text{ is even.} \end{cases} \quad (4.1.1)$$

Set also $W' = p(\widetilde{W}') \subset W$.

Assume first that $\dim V$ is even. Then $C(V)$ has a unique complex simple module S whose restriction to $C(V)_{\text{even}}$ splits into the sum of two inequivalent complex simple $C(V)_{\text{even}}$ -modules S^+, S^- . The Dirac operator $D \in \text{End}_{\mathbb{H} \otimes C(V)}(X \otimes S)$ is the endomorphism given by the action of the Dirac element \mathcal{D} . When restricted to $\mathbb{H} \otimes C(V)_{\text{even}}$, D maps $X \otimes S^{\pm}$ to $X \otimes S^{\mp}$, and denote by $D^{\pm} : X \otimes S^{\pm} \rightarrow X \otimes S^{\mp}$, the corresponding restrictions. From (3.2.3), we see that D^{\pm} both commute with the action of \widetilde{W}' . Define the Dirac cohomology of X to be the \widetilde{W} -representation

$$H_D(X) = \ker D / \ker D \cap \text{im } D. \quad (4.1.2)$$

We may also define cohomology with respect to D^{\pm} as the \widetilde{W}' -representations

$$H_D^+(X) = \ker D^+ / \ker D^+ \cap \text{im } D^- \text{ and } H_D^- = \ker D^- / \ker D^- \cap \text{im } D^+. \quad (4.1.3)$$

Then the index of X is the virtual \widetilde{W}' -module

$$I(X) = H_D^+ - H_D^-. \quad (4.1.4)$$

Now consider the case when $\dim V$ is odd. The Clifford algebra $C(V)$ has two nonisomorphic complex simple modules S^+, S^- . The restriction of S^+ and S^- to $C(V)_{\text{even}}$ are isomorphic, and they differ by the action of the center $Z(C(V)) \cong$

$\mathbb{Z}/2\mathbb{Z}$. As before, we have the Dirac operator $D \in \text{End}(X \otimes S^+)$ given by the action of \mathcal{D} . By (3.2.3), D^+ is $\text{sgn } \widetilde{W}$ -invariant, but by composing D with the vector space isomorphism $S^+ \rightarrow S^-$, we may regard $D^+ : X \otimes S^+ \rightarrow X \otimes S^-$ as \widetilde{W} -invariant. Similarly, we define D^- . Then the definitions (4.1.2), (4.1.3) and (4.1.4) make sense in this case as well. Notice that the Dirac index of $I(X)$ is a virtual $\widetilde{W}' = \widetilde{W}$ -module.

Recall that S^\pm admit structures of unitary $C(V)_{\text{even}}$ -modules. If X is unitary (or just Hermitian), let $(\ , \)_{X \otimes S^\pm}$ denote the tensor product Hermitian form on $X \otimes S^\pm$. Then D^+, D^- are adjoint with respect to $(\ , \)_{X \otimes S^\pm}$, i.e.,

$$(D^+x, y)_{X \otimes S^-} = (x, D^-y)_{X \otimes S^+}, \text{ for all } x \in X \otimes S^+, y \in X \otimes S^-. \quad (4.1.5)$$

The following result is standard.

Lemma 4.1. *As virtual \widetilde{W}' -modules, $I(X) = X \otimes S^+ - X \otimes S^-$, for every finite dimensional \mathbb{H} -module X .*

Proof. Let $\tilde{\sigma}$ be an irreducible \widetilde{W}' -module. Let $D_{\tilde{\sigma}}^\pm$ be the restrictions of D^\pm to the $\tilde{\sigma}$ -isotypic component $(X \otimes S^\pm)_{\tilde{\sigma}}$ in $X \otimes S^\pm$, respectively. There are two cases:

- (i) $D_{\tilde{\sigma}}^+ D_{\tilde{\sigma}}^- \neq 0$ or equivalently $D_{\tilde{\sigma}}^- D_{\tilde{\sigma}}^+ \neq 0$. In this case, $D_{\tilde{\sigma}}^+$ and $D_{\tilde{\sigma}}^-$ are both isomorphisms, and thus the identity in the Lemma is trivially verified.
- (ii) $D_{\tilde{\sigma}}^+ D_{\tilde{\sigma}}^- = 0$ or equivalently $D_{\tilde{\sigma}}^- D_{\tilde{\sigma}}^+ = 0$. Consider the complex

$$0 \rightarrow \ker D_{\tilde{\sigma}}^+ \rightarrow (X \otimes S^+)_{\tilde{\sigma}} \xrightarrow{D_{\tilde{\sigma}}^+} (X \otimes S^-)_{\tilde{\sigma}} \xrightarrow{D_{\tilde{\sigma}}^-} \text{im } D_{\tilde{\sigma}}^- \rightarrow 0. \quad (4.1.6)$$

The claim follows from the Euler principle. \square

Two important observations are that

$$(S^+ - S^-) \otimes (S^+ - S^-)^* \cong \frac{2}{[\widetilde{W} : \widetilde{W}']} \sum_{i=0}^n (-1)^i \wedge^i V_{\mathbb{C}}, \text{ as virtual } \widetilde{W}'\text{-modules,} \quad (4.1.7)$$

and that the character of $\sum_{i=0}^n (-1)^i \wedge^i V_{\mathbb{C}}$ on $w \in W$ is

$$\det_{V_{\mathbb{C}}}(1 - w). \quad (4.1.8)$$

In particular, since $S^+ - S^-$ is supported on the elliptic set,

$$I(\text{ind}_{\mathbb{H}_P}^{\mathbb{H}}(R_{\mathbb{C}}(\mathbb{H}_P))) = 0, \quad (4.1.9)$$

for every proper parabolic subalgebra \mathbb{H}_P of \mathbb{H} . Let $R(\widetilde{W}')_{\text{gen}}$ be the Grothendieck group of \widetilde{W}' spanned by the genuine representations, and let $\langle \ , \ \rangle_{\widetilde{W}'}$ be the usual character pairing on $R(\widetilde{W}')$.

We define an involution

$$\text{Sg} : R_{\mathbb{Z}}(\widetilde{W}') \rightarrow R_{\mathbb{Z}}(\widetilde{W}'), \quad (4.1.10)$$

as follows. If $\dim V$ is odd, $\widetilde{W}' = \widetilde{W}$, and set $\text{Sg}(\sigma) = \sigma \otimes \text{sgn}$ for every $\sigma \in R_{\mathbb{Z}}(\widetilde{W}')$. Suppose that $\dim V$ is even. Assume $\tilde{\sigma} \cong \tilde{\sigma} \otimes \text{sgn}$, for an irreducible \widetilde{W} -representation $\tilde{\sigma}$. Then $\tilde{\sigma}$ restricts to \widetilde{W}' as a sum $\tilde{\sigma}^+ \oplus \tilde{\sigma}^-$ of two inequivalent irreducible representation of the same dimension. Set $\text{Sg}(\tilde{\sigma}^\pm) = \tilde{\sigma}^\mp$. If $\tilde{\sigma} \not\cong \tilde{\sigma} \otimes \text{sgn}$, then $\tilde{\sigma}|_{\widetilde{W}'} \cong \tilde{\sigma} \otimes \text{sgn}|_{\widetilde{W}'}$, and in this case set $\text{Sg}(\tilde{\sigma}|_{\widetilde{W}'}) = \tilde{\sigma}|_{\widetilde{W}'}$. This assignment extends to an involution $\text{Sg} : R_{\mathbb{Z}}(\widetilde{W}') \rightarrow R_{\mathbb{Z}}(\widetilde{W}')$.

For every $\tilde{\sigma}_1, \tilde{\sigma}_2 \in R_{\mathbb{Z}}(\widetilde{W}')$, one has

$$\langle \tilde{\sigma}_1, \tilde{\sigma}_2 \rangle_{\widetilde{W}'} = \langle \text{Sg}(\tilde{\sigma}_1), \text{Sg}(\tilde{\sigma}_2) \rangle_{\widetilde{W}'}. \quad (4.1.11)$$

Then we obtain the following result.

Theorem 4.2 ([CT]). (1) *The map $i : R_{\mathbb{Z}}(W) \rightarrow R_{\mathbb{Z}}(\widetilde{W}')_{\text{gen}}$, $\delta \mapsto \delta \otimes (S^+ - S^-)$, gives rise to an injective map $i : \overline{R}_{\mathbb{C}}(W) \rightarrow R_{\mathbb{C}}(\widetilde{W}')_{\text{gen}}$ which satisfies*

$$\langle i(\delta_1), i(\delta_2) \rangle_{\widetilde{W}'} = 2 \langle \delta_1, \delta_2 \rangle_W^{\text{ell}}, \text{ for all } \delta_1, \delta_2 \in R_{\mathbb{C}}(W). \quad (4.1.12)$$

(2) *Let $\delta \in \overline{R}_{\mathbb{Z}}(W)$ be given. Then*

$$i(\delta) = \tilde{\delta}^+ - \tilde{\delta}^-, \quad (4.1.13)$$

for unique \widetilde{W}' -representations $\tilde{\delta}^+, \tilde{\delta}^-$ such that

$$\tilde{\delta}^- = \text{Sg}(\tilde{\delta}^+), \quad \langle \tilde{\delta}^+, \tilde{\delta}^- \rangle_{\widetilde{W}'} = 0, \text{ and } \langle \tilde{\delta}^+, \tilde{\delta}^+ \rangle_{\widetilde{W}'} = \langle \tilde{\delta}^-, \tilde{\delta}^- \rangle_{\widetilde{W}'} = \langle \delta, \delta \rangle_W^{\text{ell}}. \quad (4.1.14)$$

In particular, if δ is an element of norm one in $\overline{R}_{\mathbb{Z}}(W)$, then $\tilde{\delta}^+, \tilde{\delta}^-$ are irreducible.

Sketch of proof. We sketch the proof for convenience. Claim (1) is immediate from (4.1.7) and (4.1.8). For claim (2), one uses the involution Sg from (4.1.10). Notice that

$$\text{Sg}(i(\delta)) = -i(\delta).$$

Since $i(\delta)$ is an integral virtual \widetilde{W}' -character, there exist \widetilde{W}' -representations $\tilde{\delta}^+$ and $\tilde{\delta}^-$ with $\langle \tilde{\delta}^+, \tilde{\delta}^- \rangle_{\widetilde{W}'} = 0$, such that $i(\delta) = \tilde{\delta}^+ - \tilde{\delta}^-$. Applying Sg , we conclude that $\tilde{\delta}^- = \text{Sg}(\tilde{\delta}^+)$. The rest is a consequence of (1). \square

Motivated by this result, we make the following definition.

Definition 4.3. Let $\text{Irr}_{\text{gen}}^0 \widetilde{W}'$ be the set of irreducible genuine \widetilde{W}' -representations $\tilde{\delta}$ which occur as a component of $i(\delta)$ for some $\delta \in \overline{R}_{\mathbb{Z}}(W)$, see Theorem 4.2(2). From the construction, one sees that if $\tilde{\delta} \in \text{Irr}_{\text{gen}}^0 \widetilde{W}'$, then $\text{Sg}(\tilde{\delta}) \in \text{Irr}_{\text{gen}}^0 \widetilde{W}'$.

Example 4.4. If R is of type A_{n-1} , then $\text{Irr}_{\text{gen}}^0 \widetilde{S}'_n = \{S^+, S^-\}$, where S^+, S^- are the two associate spin \widetilde{S}'_n -modules. For comparison, in this case, $|\text{Irr}_{\text{gen}} \widetilde{S}'_n| = k_1 + 2k_2$, where k_1 is the number of partitions of n of even length, and k_2 is the number of partitions of n of odd length.

If R is of type B_n , then $\text{Irr}_{\text{gen}}^0 \widetilde{W}' = \text{Irr}_{\text{gen}} \widetilde{W}'$. This can be seen directly using the description of $\widetilde{W}(B_n)$ -representations, see [C, Theorem 1.0.1 and section 3.3].

4.2. The central character and Vogan's conjecture. Theorem 3.2 imposes strict limitations on the central character of modules with nonzero Dirac cohomology. In light of Corollary 3.3, we may regard the homomorphism ζ as

$$\zeta : Z(\mathbb{H}) \rightarrow Z(\mathbb{C}[\widetilde{W}']).$$

Definition 4.5. If $\tilde{\delta}$ is an irreducible \widetilde{W}' -representation, define the homomorphism (central character) $\chi^{\tilde{\delta}} : Z(\mathbb{H}) \rightarrow \mathbb{C}$ by $\chi^{\tilde{\delta}}(z) = \tilde{\delta}(\zeta(z))$ for every $z \in Z(\mathbb{H})$.

As in [BCT], an immediate corollary of Theorem 3.2 is the following.

Corollary 4.6. *Let X be a \mathbb{H} -module with central character χ_X . Suppose there exists an irreducible \widetilde{W}' -representation $\widetilde{\delta}$ such that $\text{Hom}_{\widetilde{W}'}[\widetilde{\delta}, H_D(X)] \neq 0$. Then $\chi_X = \chi^{\widetilde{\delta}}$.*

4.3. The rank pairing. Let $K_0(\mathbb{H})$ be the Grothendieck group of finitely generated projective \mathbb{H} -modules. Every such projective module P is given by an idempotent $p \in M_N(\mathbb{H})$, for some natural number N , such that P can be identified with the image of p acting on \mathbb{H}^N . One defines the rank pairing to be the bilinear map

$$[\cdot, \cdot] : K_0(\mathbb{H}) \times R_{\mathbb{Z}}(\mathbb{H}) \rightarrow \mathbb{Z}, \quad \text{such that } [[P], [\pi]] := \text{rank}(\pi(p)), \quad (4.3.1)$$

for every finitely generated projective module P and every irreducible \mathbb{H} -module (π, V_π) ; here $\pi(p)$ is regarded as an element of $M_N(\text{End}_{\mathbb{H}}(V_\pi))$.

The rank pairing extends to a Hermitian pairing $[\cdot, \cdot] : K_0(\mathbb{H})_{\mathbb{C}} \times R_{\mathbb{C}}(\mathbb{H}) \rightarrow \mathbb{C}$.

Remark 4.7. If δ is a finite dimensional W -representation, the induced module $\mathbb{H} \otimes_{\mathbb{C}[W]} \delta$, an \mathbb{H} -module under left multiplication, is a finitely generated projective \mathbb{H} -module. For every finite dimensional \mathbb{H} -module X , the rank pairing is

$$[\mathbb{H} \otimes_{\mathbb{C}[W]} \delta, X] = \dim \text{Hom}_W(\delta, X). \quad (4.3.2)$$

The same formula holds with W' in place of W .

We define a notion of support for elements of $K_0(\mathbb{H})_{\mathbb{C}}$. If $\lambda \in V_{\mathbb{C}}^{\vee}$, let $\text{Irr}_{W\lambda}(\mathbb{H})$ denote the set of isomorphism classes of irreducible \mathbb{H} -modules with central character $W\lambda$.

Definition 4.8. If $A \in K_0(\mathbb{H})_{\mathbb{C}}$, define

$$\text{supp}(A) = \{\lambda \in V_{\mathbb{C}}^{\vee} : \text{there exists } X \in \text{Irr}_{W\lambda}(\mathbb{H}) \text{ such that } [A, X] \neq 0\}. \quad (4.3.3)$$

Set $F_i K_0(\mathbb{H})_{\mathbb{C}} = \{A \in K_0(\mathbb{H})_{\mathbb{C}} : \dim \text{supp}(A) \leq i\}$ and $F_{-1} K_0(\mathbb{H})_{\mathbb{C}} = \cap_{\pi \in \text{Irr}(\mathbb{H})} \ker[\cdot, \pi]$. Define

$$F_0 H_0(\mathbb{H}) = F_0 K_0(\mathbb{H})_{\mathbb{C}} / F_{-1} K_0(\mathbb{H})_{\mathbb{C}}.$$

By definition,

$$[A, X] = 0 \text{ for all } X \in \text{Irr}(\mathbb{H}) \text{ implies that } A \in F_{-1} K_0(\mathbb{H})_{\mathbb{C}}. \quad (4.3.4)$$

Lemma 4.9. *The rank pairing (4.3.1) gives rise to a canonical injective linear map $\Phi : F_0 H_0(\mathbb{H}) \rightarrow \overline{R}_{\mathbb{C}}(\mathbb{H})$ such that*

$$[A, X] = \langle \Phi(A), X \rangle_{\mathbb{H}}^{\text{EP}}, \text{ for all } A \in F_0 H_0(\mathbb{H}), X \in \overline{R}_{\mathbb{C}}(\mathbb{H}). \quad (4.3.5)$$

Proof. Firstly, the rank pairing descends to a pairing $[\cdot, \cdot] : F_0 H_0(\mathbb{H}) \times R_{\mathbb{C}}(\mathbb{H}) \rightarrow \mathbb{C}$, which is nondegenerate on the left by (4.3.4). Secondly, let $A \in F_0 H_0(\mathbb{H})$ be given, and suppose that $X = \mathbb{H} \otimes_{\mathbb{H}_P} \sigma$ is a parabolically induced module, $\sigma \in \text{Irr}(\mathbb{H}_P)$ such that $[A, X] \neq 0$. We can form the family $\{\sigma_\chi = \sigma \otimes \chi : \chi \text{ character of } Z(\mathbb{H}_P)\} \subset \text{Irr}(\mathbb{H}_P)$. Then $[A, X_\chi] \neq 0$, for all $X_\chi = \mathbb{H} \otimes_{\mathbb{H}_P} \sigma_\chi$. But, by definition, this implies that the central characters of X_χ are all in $\text{supp}(A)$, and therefore $\dim \text{supp}(A) > 0$, a contradiction. Hence, $[A, X] = 0$ for every proper parabolically induced module X .

This implies that the rank pairing descends to a pairing $F_0 H_0(\mathbb{H}) \otimes \overline{R}_{\mathbb{C}}(\mathbb{H}) \rightarrow \mathbb{C}$. Because this is nondegenerate on the left, it defines an injective linear map

$$\Phi' : F_0 H_0(\mathbb{H}) \rightarrow \overline{R}_{\mathbb{C}}(\mathbb{H})^*, \quad \Phi'(A) = [A, \cdot] : \overline{R}_{\mathbb{C}}(\mathbb{H}) \rightarrow \mathbb{C}. \quad (4.3.6)$$

Using the nondegenerate Hermitian pairing $\langle \cdot, \cdot \rangle_{\mathbb{H}}^{\text{EP}} : \overline{R}_{\mathbb{C}}(\mathbb{H}) \times \overline{R}_{\mathbb{C}}(\mathbb{H}) \rightarrow \mathbb{C}$, we get an injection

$$\iota : \overline{R}_{\mathbb{C}}(\mathbb{H}) \rightarrow \overline{R}_{\mathbb{C}}(\mathbb{H})^*, \quad \iota(X) = \langle \cdot, X \rangle_{\mathbb{H}}^{\text{EP}}, \quad X \in \overline{R}_{\mathbb{C}}(\mathbb{H}). \quad (4.3.7)$$

Since $\overline{R}_{\mathbb{C}}(\mathbb{H})$ is finite dimensional, ι is a bijection. Define

$$\Phi = \iota^{-1} \circ \Phi' : F_0 H_0(\mathbb{H}) \rightarrow \overline{R}_{\mathbb{C}}(\mathbb{H}). \quad (4.3.8)$$

It is clear from the definitions that $[A, X] = \langle \Phi(A), X \rangle_{\mathbb{H}}^{\text{EP}}$, for all $A \in F_0 H_0(\mathbb{H})$, $X \in \overline{R}_{\mathbb{C}}(\mathbb{H})$. \square

4.4. The Dirac induction map Ind_D .

Definition 4.10. The Dirac induction map $\text{Ind}_D : \overline{R}_{\mathbb{C}}(W) \rightarrow \overline{R}_{\mathbb{C}}(\mathbb{H})$ is the linear map defined on every $\delta \in \overline{R}_{\mathbb{Z}}(W)$ by

$$\text{Ind}_D(\delta) = \Phi([\mathbb{H} \otimes_{W'} ((\tilde{\delta}^+)^* \otimes S^+)] - [\mathbb{H} \otimes_{W'} ((\tilde{\delta}^+)^* \otimes S^-)]), \quad (4.4.1)$$

where Φ is the map from Lemma 4.9 and δ^+ is the \widetilde{W}' -representation from Theorem 4.2(2).

Theorem 4.11. (1) *The map Ind_D from Definition 4.10 is well-defined, i.e.,*
 $[\mathbb{H} \otimes_{W'} ((\tilde{\delta}^+)^* \otimes S^+)] - [\mathbb{H} \otimes_{W'} ((\tilde{\delta}^+)^* \otimes S^-)] \in F_0 H_0(\mathbb{H}).$
 (2) *For every $\delta \in \overline{R}_{\mathbb{Z}}(W)$,*

$$\langle \text{Ind}_D(\delta), X \rangle_{\mathbb{H}}^{\text{EP}} = \langle \tilde{\delta}^+, I(X) \rangle_{\widetilde{W}'}, \quad (4.4.2)$$

where $I(X)$ is the Dirac index of X .

(3) *For every $X \in \overline{R}_{\mathbb{C}}(\mathbb{H})$ and $\delta \in \overline{R}_{\mathbb{C}}(W)$,*

$$\langle \text{Ind}_D(\delta), X \rangle_{\mathbb{H}}^{\text{EP}} = \langle \delta, r(X) \rangle_W^{\text{ell}}. \quad (4.4.3)$$

Proof. Let X be a finite dimensional \mathbb{H} -module. Let δ be an element of $\overline{R}_{\mathbb{Z}}(\mathbb{H})$. Using (4.3.2), we have the rank pairings:

$$[\mathbb{H} \otimes_{W'} ((\tilde{\delta}^+)^* \otimes S^{\pm}), X] = \dim \text{Hom}_{W'}((\tilde{\delta}^+)^* \otimes S^{\pm}, X) = \dim \text{Hom}_{\widetilde{W}'}((\delta^+)^*, X \otimes (S^{\pm})^*).$$

Therefore, using also Lemma 4.1:

$$\begin{aligned} [\mathbb{H} \otimes_{W'} ((\tilde{\delta}^+)^* \otimes S^+), X] - [\mathbb{H} \otimes_{W'} ((\tilde{\delta}^+)^* \otimes S^-), X] &= \dim \text{Hom}_{\widetilde{W}'}((\delta^+)^*, I(X)^*) \\ &= \langle \tilde{\delta}^+, I(X) \rangle_{\widetilde{W}'}. \end{aligned} \quad (4.4.4)$$

Since $I(X) = 0$ whenever X is a proper parabolically induced module (4.1.9), we see that the support of $[\mathbb{H} \otimes_{W'} ((\tilde{\delta}^+)^* \otimes S^+), X] - [\mathbb{H} \otimes_{W'} ((\tilde{\delta}^+)^* \otimes S^-), X]$ is a subset of the set of central characters of elliptic tempered \mathbb{H} -modules. Then (1) follows.

Claim (2) is immediate from (4.4.4) and Lemma 4.9.

For claim (3) let δ be an element of $\overline{R}_{\mathbb{Z}}(W)$ and X an element of $\overline{R}_{\mathbb{Z}}(\mathbb{H})$. Using Theorem 4.2(1),

$$\begin{aligned} \langle \delta, X|_W \rangle_W^{\text{ell}} &= \frac{1}{2} \langle i(\delta), i(X|_W) \rangle_{\widetilde{W}'} = \frac{1}{2} \langle i(\delta), I(X) \rangle_{\widetilde{W}'}, \\ &= \frac{1}{2} \langle \tilde{\delta}^+ - \tilde{\delta}^-, I(X) \rangle_{\widetilde{W}'} = \langle \tilde{\delta}^+, I(X) \rangle_{\widetilde{W}'}. \end{aligned} \quad (4.4.5)$$

For the last equality, we used the involution Sg of (4.1.10) to see that

$$\langle \tilde{\delta}^+, I(X) \rangle_{\widetilde{W}'} = \langle \text{Sg}(\tilde{\delta}^+), \text{Sg}(I(X)) \rangle_{\widetilde{W}'} = \langle \tilde{\delta}^-, -I(X) \rangle_{\widetilde{W}'}.$$

□

Definition 4.12. Let $(\mathcal{X}, \langle \cdot, \cdot \rangle)$ be a \mathbb{Z} -lattice in an Euclidean space $(E, \langle \cdot, \cdot \rangle)$. A vector $v \in \mathcal{X}$ is called pure if v cannot be written as a sum $v = v_1 + v_2$, where $v_1, v_2 \in \mathcal{X} \setminus \{0\}$ and $\langle v_1, v_2 \rangle = 0$.

Denote

$$\mathcal{Y} = \mathbf{r}(\overline{R}_{\mathbb{Z}}(\mathbb{H})) \subset \overline{R}_{\mathbb{Z}}(W), \quad (4.4.6)$$

a \mathbb{Z} -lattice with respect to the pairing $\langle \cdot, \cdot \rangle_W^{\text{ell}}$ in $\overline{R}_{\mathbb{C}}(W)$. Notice that a priori, the lattice \mathcal{Y} depends on the parameter function k of \mathbb{H} .

Corollary 4.13. (1) Ind_D is the inverse map of \mathbf{r} .

(2) For every $\delta \in \overline{R}_{\mathbb{C}}(W)$, the Dirac index of $\text{Ind}_D(\delta)$ is

$$I(\text{Ind}_D(\delta)) = i(\delta). \quad (4.4.7)$$

In particular, $I(\text{Ind}_D(\delta))$ is independent of the parameter function k of \mathbb{H} .

(3) If δ is a pure element (in the sense of Definition 4.12) of the lattice \mathcal{Y} defined in (4.4.6), then $\text{Ind}_D(\delta)$ is supported by a single central character.

(4) If a rational multiple of $\delta \in \overline{R}_{\mathbb{Z}}(W)$ is pure in \mathcal{Y} , then the central character of $\text{Ind}_D(\delta)$ equals $\chi^{\tilde{\delta}}$ (see Definition 4.5) for any irreducible \widetilde{W}' -representation $\tilde{\delta}$ occurring in $i(\delta)$.

Proof. Claim (1) is immediate by Theorem 4.11(3). Namely, given $X \in \overline{R}_{\mathbb{C}}(\mathbb{H})$:

$$\langle \text{Ind}_D(\mathbf{r}(X)), Y \rangle_{\mathbb{H}}^{\text{EP}} = \langle \mathbf{r}(X), \mathbf{r}(Y) \rangle_W^{\text{ell}} = \langle X, Y \rangle_{\mathbb{H}}^{\text{EP}}, \quad (4.4.8)$$

for every $Y \in \overline{R}_{\mathbb{C}}(\mathbb{H})$. Since $\langle \cdot, \cdot \rangle_{\mathbb{H}}^{\text{EP}}$ is nondegenerate on $\overline{R}_{\mathbb{C}}(\mathbb{H})$, it follows that $\text{Ind}_D(\mathbf{r}(X)) = X$ in $\overline{R}_{\mathbb{C}}(\mathbb{H})$.

By Lemma 4.1, $I(\text{Ind}_D(\delta)) = i(\mathbf{r}(\text{Ind}_D(\delta)))$. Claim (2) now follows from (1).

For (3), the restriction $\text{Ind}_D|_{\mathcal{Y}} : \mathcal{Y} \rightarrow \overline{R}_{\mathbb{Z}}(\mathbb{H})$ is an isometric isomorphism and an inverse of $\mathbf{r} : \overline{R}_{\mathbb{Z}}(\mathbb{H}) \rightarrow \mathcal{Y}$. Given $\delta \in \mathcal{Y}$, decompose $\text{Ind}_D(\delta) = X_1 + \cdots + X_{\ell}$ in $\overline{R}_{\mathbb{Z}}(H)$, where:

- (a) X_i is in the class of an integral virtual \mathbb{H} -module;
- (b) X_i has central character χ_i ;
- (c) the central characters χ_i are mutually distinct.

By (c), $\text{EP}(X_i, X_j) = 0$ for all $i \neq j$. Then $\delta = \mathbf{r}(\text{Ind}_D(\delta)) = \mathbf{r}(X_1) + \cdots + \mathbf{r}(X_{\ell})$ is an orthogonal decomposition of δ in \mathcal{Y} . Since δ is pure, $\ell = 1$ by definition.

To prove (4), notice that (3) implies that $\text{Ind}_D(\delta)$ is supported by a single central character. On the other hand, if $\tilde{\delta}$ occurs in $i(\delta)$, then $\langle \tilde{\delta}, I(\text{Ind}_D(\delta)) \rangle_{\widetilde{W}'} \neq 0$. This means that $\tilde{\delta}$ occurs in (one of) the Dirac cohomology groups $H_D(\text{Ind}_D(\delta))$, so the claim follows from Corollary 4.6. □

Remark 4.14. (a) Corollary 4.13(3) says in particular that if a rational multiple of $\delta \in \overline{R}_{\mathbb{Z}}(W)$ is pure in \mathcal{Y} , then for all \widetilde{W}' -irreducible constituents $\tilde{\delta}$ of $i(\delta)$, the central characters $\chi^{\tilde{\delta}}$ are the same. Define the central character of Ind_D to be

$$\text{cc}(\text{Ind}_D(\delta)) = \chi^{\tilde{\delta}}, \quad (4.4.9)$$

for every $\delta \in \overline{R}_{\mathbb{Z}}(W)$ which is a rational multiple of a pure element in \mathcal{Y} .

- (b) Every Euclidean lattice L has a basis consisting of pure vectors. Recall that if \mathcal{B} is a basis for L , then the orthogonal defect of \mathcal{B} is defined as the ratio $\text{def}(\mathcal{B}) = \prod_{x \in \mathcal{B}} \|x\| / \text{vol}(L)$. Then every basis with minimal orthogonal defect consists of pure vectors.

In our situation, this implies that \mathcal{Y} has a basis of pure elements, and therefore that every element of $\overline{R}_{\mathbb{Z}}(W)$ can be written as a sum of elements of $\overline{R}_{\mathbb{Z}}(W)$ which are rational multiples of pure vectors in \mathcal{Y} .

- (c) We will improve Corollary 4.13(4) in Corollary 5.7.

4.5. The central character map. Let \mathbb{H}_A be the generic Hecke algebra over $A = \mathbb{C}[\underline{k}]$ from Definition 2.2.

Definition 4.15. Let $\text{Res}^{\text{lin}}(R)$ be the set of linear maps $\xi : \text{Spec}(A) = \mathbb{C}^{W \setminus R} \rightarrow V_{\mathbb{C}}^{\vee}$ such that for almost all $k \in \mathbb{C}^{W \setminus R}$, the point $\xi(k)$ satisfies the condition

$$\#\{\alpha \in R : \alpha(\xi(k)) = k_{\alpha}\} = \#\{\alpha \in R : \alpha(\xi(k)) = 0\} + \dim V_{\mathbb{C}}^{\vee}. \quad (4.5.1)$$

The set $\text{Res}^{\text{lin}}(R)$ is studied in [HO, O1, O2], where it is shown that $\text{Res}^{\text{lin}}(R)$ is a nonempty, finite set, invariant under W , and its explicit description is given in all irreducible cases. Notice that when k is a constant function, condition (4.5.1) is satisfied by the middle elements of distinguished Lie triples [Ca].

We will make repeated use of the following results. Assume the root system R does not have simply-laced factors and recall the notion of generic parameter k from [O1, section 4.3] and [OS2, Definition 2.64].

Theorem 4.16 ([OS2, Theorem 5.3, Definition 5.4, equation (79)]). *Suppose R does not have simply laced factors, and let \mathcal{Q} be a generic region for the parameters k .*

- (1) *If π is an irreducible discrete series \mathbb{H}_k -module, there exists $\lambda \in \text{Res}^{\text{lin}}(R)$ such that the central character of π is $\lambda(k)$.*
- (2) *Conversely, if $\lambda \in \text{Res}^{\text{lin}}(R)$, for every $k \in \mathcal{Q}$ there exists a unique irreducible discrete series \mathbb{H}_k -module π_k with central character $\lambda(k)$.*

Combining Theorem 4.16 with the results of [OS1] on the Euler-Poincaré pairing as recalled in sections 2.5 and 2.6, one has the following result.

Theorem 4.17 ([OS1, OS2]). *Suppose R does not have simple laced factors, and k is a generic parameter for \mathbb{H} . Then the set of irreducible discrete series modules $\text{DS}(\mathbb{H})$ is an orthonormal basis for $\overline{R}_{\mathbb{Z}}(\mathbb{H})$.*

Motivated by the results of the previous section, let $\mathcal{S}(\overline{R}_{\mathbb{Z}}(W), \mathcal{Y})$ be the set of elements $\delta \in \overline{R}_{\mathbb{Z}}(W)$ which are rational multiples of pure elements of \mathcal{Y} . Define

$$\Lambda : \mathcal{S}(\overline{R}_{\mathbb{Z}}(W), \mathcal{Y}) \rightarrow \text{Spec}(Z(\mathbb{H}_A)), \quad \Lambda(\delta)(k) = \text{cc}(\text{Ind}_{D,k}(\delta)) \in \text{Spec}(Z(\mathbb{H}_k)) = W \setminus V_{\mathbb{C}}^{\vee}. \quad (4.5.2)$$

Using Definition 4.3 and Corollary 4.13(4), we see that

$$\text{im } \Lambda = \{\chi^{\tilde{\delta}} : \tilde{\delta} \in \text{Irr}_{\text{gen}}^0(\widetilde{W}')\}. \quad (4.5.3)$$

- Theorem 4.18.** (1) *Let $\tilde{\delta} \in \text{Irr}_{\text{gen}}^0(\widetilde{W}')$ be given. Then $\chi^{\tilde{\delta}}$ is linear in \underline{k} , i.e., there exists a linear function $\xi(\tilde{\delta}) : \text{Spec}(A) \rightarrow V_{\mathbb{C}}^{\vee}$ such that $\chi^{\tilde{\delta}} = W \cdot \xi(\tilde{\delta})$.*
- (2) *$\text{Res}^{\text{lin}}(R)$ is contained in $\text{im } \Lambda$, with equality if R has no simply-laced factors.*

Proof. (1) Let $\delta \in \overline{R}_{\mathbb{Z}}(W)$ be such that $\tilde{\delta}$ is a component of $i(\delta)$. Since \mathcal{Y} admits a basis of pure vectors, we may assume that δ is pure. The equality $\Lambda(\delta) = \chi^{\tilde{\delta}}$ was verified in Corollary 4.13(4). It is sufficient to prove the linearity of $\Lambda(\delta)$ in the case when R is irreducible.

Assume that R is simply laced. The generic algebra \mathbb{H}_A admits scaling isomorphisms $s_c : \mathbb{H}_k \rightarrow \mathbb{H}_{ck}$, for a scalar c , given by $s_c(w) = w$, $w \in W$ and $s_c(v) = cv$, for $v \in V$. Notice that the Dirac element \mathcal{D} is unchanged under s_c . For a central element $z \in S(V_{\mathbb{C}})^W$ homogeneous of degree N , equation (3.3.1) shows that $\zeta_{ck}(z)$ has degree N in c . Therefore,

$$\zeta_{ck}(z) = c^N \zeta_k(z), \text{ and in particular } \zeta_k(z) = k^N \zeta_1(z).$$

Hence $\chi^{\tilde{\delta}}$ is linear in k for all irreducible \widetilde{W}' -representations $\tilde{\delta}$.

Now, let R be irreducible, non-simply laced. Fix a generic region \mathcal{Q} of the parameter function k , and $k_0 \in \mathcal{Q}$. Corollary 4.13(4) and Theorem 4.16 imply that there exists $\lambda \in \text{Res}^{\text{lin}}(R)$ such that $\Lambda(\delta)(k) = \lambda(k)$ for every $k \in \mathcal{Q}$. Consider the formal completion $\widehat{\mathbb{C}[k]}$ of $\mathbb{C}[k]$ at k_0 , and $i_{k_0} : \mathbb{C}[k] \rightarrow \widehat{\mathbb{C}[k]}$ the canonical map. Since

$$\Lambda(\delta) \circ i_{k_0} = \lambda \circ i_{k_0},$$

as homomorphisms $S(V_{\mathbb{C}})^W \rightarrow \widehat{\mathbb{C}[k]}$, it follows from the injectivity of i_{k_0} that $\Lambda(\delta) = \lambda$.

(2) When k is constant (more generally when k is of geometric origin in the sense of [L2]), this is (part of) [BCT, Theorem 5.8], which is a corollary of Vogan's conjecture 3.2 together with [C, Theorems 1.0.1 and 3.10.3]. In particular, this is the case when R is simply-laced. The explicit map Λ for these cases can be found in the tables of [C].

Assume now that R is a simple non-simply laced root system. By (1), $\Lambda(\delta)$ gives a W -orbit of linear maps $\mathbb{C}^{(W \setminus R)} \rightarrow V_{\mathbb{C}}^{\vee}$. So it remains to check condition (4.5.1) in one generic region of k . To simplify notation, let k be the parameter on the long roots of R and k' the parameter on the short roots.

When R is of type B_n , we choose the generic region $k'/k > n - 1$. To every partition σ of n , [O1] attaches an element cc_{σ} of $\text{Res}^{\text{lin}}(R)$, and every W -conjugacy class in $\text{Res}^{\text{lin}}(R)$ contains one and only one such cc_{σ} . Let $\sigma \times \emptyset$ denote the irreducible $W(B_n)$ -representation obtained by inflating to $W(B_n)$ the irreducible S_n -representation given in Young's parametrization by σ . An easy algebraic argument (see [CK, section 4.7]) shows that there is a unique discrete series π_{σ} with central character cc_{σ} and $\pi_{\sigma}|_W \cong (\sigma \times \emptyset) \otimes \text{sgn}$. It is well-known (see [Re]) that $(\sigma \times \emptyset) \otimes S$ is an irreducible \widetilde{W} -representation, for each spin \widetilde{W} -module S , and in fact, every genuine irreducible \widetilde{W} -representation is obtained in this way. Fix a spin module S and denote $\tilde{\sigma} = (\sigma \times \emptyset) \otimes S \otimes \text{sgn}$. Then

$$I(\pi_{\sigma}) = \pi_{\sigma}|_W \otimes (S^+ - S^-) = (\sigma \times \emptyset) \otimes (S^+ - S^-) \otimes \text{sgn} \neq 0. \quad (4.5.4)$$

Therefore, $\text{Hom}_{\widetilde{W}}[\tilde{\sigma}, H_D(\pi_{\sigma})] \neq 0$, and by Corollary 4.6, $\chi^{\tilde{\sigma}} = \text{cc}_{\sigma}$. This completes the proof in type B_n .

For types G_2 and F_4 , the notation for genuine \widetilde{W} -types is as in the character tables of [M]. By (3.3.2), if $\Lambda(\delta) = \xi \in \text{Res}^{\text{lin}}(R)$, then

$$\langle \xi(k, k'), \xi(k, k') \rangle = \tilde{\delta}(\Omega_{\widetilde{W}, k, k'}),$$

as functions in k, k' . It turns out that for F_4 and G_2 , if $\xi \neq \xi' \in W \setminus \text{Res}^{\text{lin}}(R)$, then $\langle \xi(k, k'), \xi(k, k') \rangle \neq \langle \xi'(k, k'), \xi'(k, k') \rangle$ as functions in k, k' . (This is not the case for type B_n discussed above, when n is large.) Therefore, to identify the images of the map Λ , it is sufficient in this case to compute the scalar functions $\tilde{\delta}(\Omega_{\widetilde{W}, k, k'})$ and compare. The results are tabulated in Tables 1 and 2.

TABLE 1. $\text{Res}^{\text{lin}}(F_4)$

central character	$\tilde{\delta} \in \text{Irr}_{\text{gen}}^0(\widetilde{W})/\sim$
$k\omega_1 + k\omega_2 + k'\omega_3 + k'\omega_4$	4_s
$k\omega_1 + k\omega_2 + (-k + k')\omega_3 + k'\omega_4$	8_{sss}
$k\omega_1 + k\omega_2 + (-k + k')\omega_3 + k\omega_4$	12_s
$k\omega_1 + k\omega_2 + (-2k + k')\omega_3 + k'\omega_4$	4_{ss}
$k\omega_1 + k\omega_2 + (-2k + k')\omega_3 + 2k\omega_4$	24_s
$k\omega_1 + k\omega_2 + (-2k + k')\omega_3 + k\omega_4$	12_{ss}
$k\omega_1 + k\omega_2 + (-2k + k')\omega_3 + (3k - k')\omega_4$	8_{ssss}
$k\omega_2 + (-k + k')\omega_4$	$8_{ss}, 8_s$

TABLE 2. $\text{Res}^{\text{lin}}(G_2)$

central character	$\tilde{\delta} \in \text{Irr}_{\text{gen}}^0(\widetilde{W})/\sim$
$k\omega_1 + k'\omega_2$	2_s
$k\omega_1 + (-k + k')\omega_2$	2_{ss}
$k\omega_1 + \frac{1}{2}(-k + k')$	2_{sss}

□

5. ORTHOGONAL BASES FOR SPACES OF VIRTUAL ELLIPTIC CHARACTERS

We would like to describe the map Ind_D from Definition 4.10 explicitly and study its integrality properties. For this, we will show that the lattice $\overline{R}_{\mathbb{Z}}(W)$ admits an orthogonal basis for all irreducible root systems R . In particular, this basis consists of pure vectors, in the sense of Definition 4.12.

In Theorem 5.1, we determine orthogonal bases for the spaces of virtual elliptic characters $\overline{R}_{\mathbb{Z}}(\mathbb{H})$, for every irreducible root system R and every parameter function k , except when $R = F_4$ and k is special nonconstant. By Corollary 2.11, we see that every orthogonal basis of $\overline{R}_{\mathbb{Z}}(\mathbb{H})$ with respect to $\langle \cdot, \cdot \rangle_{\mathbb{H}}^{\text{EP}}$ gives, by restriction to W , an orthogonal basis of $\overline{R}_{\mathbb{Z}}(W)$ with respect to $\langle \cdot, \cdot \rangle_W^{\text{ell}}$.

When R is non-simply laced, Theorem 4.17 gives a complete answer for generic parameter functions. For special parameters (i.e., non-generic) we use a limiting argument. This argument is available when R is of type B_n via the exotic geometric models for \mathbb{H} -modules of [K], and it was already used in [CKK]. This approach can also be applied when R is of type D_n , since the graded Hecke algebra of type B_n with parameter 0 on the short roots is a $\mathbb{Z}/2\mathbb{Z}$ extension of the graded Hecke algebra of type D_n . However, in the D_n case, one needs to analyze carefully the changes in the R -groups under the extension by $\mathbb{Z}/2\mathbb{Z}$, using the explicit description of type B_n R -groups in [Sl]. The same limit argument can be used for $R = G_2$, where it

easy to construct explicit models for the families of discrete series in the generic regions. Thus the cases that we cannot treat here are certain special values of the parameters when $R = F_4$.

When R is simply laced, especially if R is of type E , we need to use the geometric classification of [KL] and the results in [R] relating the elliptic theories with the geometry of Kazhdan and Lusztig. Let \mathfrak{g} be the complex simple Lie algebra with Cartan subalgebra $V_{\mathbb{C}}^{\vee}$ and root system R and G be the complex connected adjoint Lie group with algebra \mathfrak{g} . By [KL, L1, L2], when the parameter function is constant $k = 1$, the irreducible \mathbb{H} -modules are parametrized by G -conjugacy classes of triples (s, e, ψ) , where $s \in \mathfrak{g}$ is semisimple, $e \in \mathfrak{g}$ is such that $[s, e] = e$, in particular, e is nilpotent, and ψ is an irreducible representation of Springer type of the group of components $A(s, e)$ of the centralizer in G of s and e . Write $\pi_{(s, e, \psi)}$ for the module parametrized by the class $[(s, e, \psi)]$. In this correspondence, tempered modules with real central character are attached to triples $(\frac{1}{2}h, e, \psi)$, for a Lie triple (e, h, f) , while discrete series are attached to $(\frac{1}{2}h, e, \psi)$ where e is distinguished in the sense of Bala-Carter [Ca]. Thus tempered modules with real central character are uniquely determined by e and $\psi \in \hat{A}(e)_0$ (characters of the component group of Springer type), so we write $\pi_{e, \psi}$ in place of $\pi_{(\frac{1}{2}h, e, \psi)}$.

Reeder's results [R] imply in this case that an irreducible tempered module $\pi_{(e, \psi)} \equiv 0$ in $\overline{R}_{\mathbb{Z}}(\mathbb{H})$, unless e is a quasidistinguished nilpotent element (see [R, (3.2.2)] for the definition, recalled below). If the parameter function k is constant $k = 1$, define the following set, consisting of elliptic tempered modules,

$$\mathcal{B}(\overline{R}_{\mathbb{Z}}(\mathbb{H})) = \{\pi_{e, \psi} : e \text{ dist.}, \psi \in \hat{A}(e)_0\} \cup \{\pi_{e, \text{triv}} : e \text{ quasidist., not dist.}\}. \quad (5.0.5)$$

The notation for nilpotent orbits below is as in [Ca].

- Theorem 5.1.** (1) *Assume the root system R is not of type D_{2n} or E_7 , and if $R = F_4$ then the parameter function k is assumed either constant or generic. The space $\overline{R}_{\mathbb{Z}}(\mathbb{H})$ has an orthonormal basis with respect to $\langle \cdot, \cdot \rangle_{\mathbb{H}}^{\text{EP}}$ consisting of elliptic tempered \mathbb{H} -representations. In particular, when the parameter function is constant $k = 1$, the set $\mathcal{B}(\overline{R}_{\mathbb{Z}}(\mathbb{H}))$ from (5.0.5) is such an orthonormal basis.*
- (2) *Suppose $R = D_{2n}$. The set $\mathcal{B}(\overline{R}_{\mathbb{Z}}(\mathbb{H}))$ is an orthogonal basis of $\overline{R}_{\mathbb{Z}}(\mathbb{H})$ such that every element is a unit element, except the irreducible tempered modules $\pi_{e, \text{triv}}$ when e are representatives of quasidistinguished nilpotent orbits labeled by partitions $(a_1, a_1, a_2, a_2, \dots, a_{2l}, a_{2l})$ of $2n$, $0 < a_1 < a_2 < \dots < a_{2l}$, have elliptic norm $\sqrt{2}$.*
- (3) *Suppose $R = E_7$. The set $\mathcal{B}(\overline{R}_{\mathbb{Z}}(\mathbb{H}))$ is an orthogonal basis for $\overline{R}_{\mathbb{Z}}(\mathbb{H})$ such that every element is a unit element, except the irreducible tempered module $\pi_{A_4+A_1, \text{triv}}$ has elliptic norm $\sqrt{2}$.*

The proof of Theorem 5.1 is case by case and it will occupy the rest of the section. The case $R = A_{n-1}$ is well-known: the space $\overline{R}_{\mathbb{Z}}(\mathbb{H})$ is one-dimensional spanned by the Steinberg module.

5.1. Non-simply laced R . When the parameter function k is generic, Theorem 4.17 says that $\text{DS}(\mathbb{H})$ is an orthonormal basis of $\overline{R}_{\mathbb{Z}}(\mathbb{H})$, and therefore this proves Theorem 5.1(1) in this case.

When k is non-generic, one proves the result by a known limiting argument, e.g., [CK, section 2.4]. Assume R is of type B_n or G_2 . Let k_0 be a special parameter

function. Without loss of generality, we may assume that $k_0(\alpha) = 1$, when α is a long root, and $k_0(\beta) = m_0$, when β is a short root. Then there exists $\epsilon > 0$ such that the parameter function k_t , where $k_t(\alpha) = 1$ and $k_t(\beta) = m_0 + t$ is generic for all $t \in (0, \epsilon)$. Let

$$\mathcal{F} = \{\pi_t^{\mathcal{F}} : \pi_t^{\mathcal{F}} \in \text{DS}(\mathbb{H}_{k_t})\}_{0 < t < \epsilon}$$

be a continuous family of discrete series as in [OS2, Definition 3.5]. Then $\pi_t^{\mathcal{F}}|_W \cong \pi_{t'}^{\mathcal{F}}|_W$ for all $t, t' \in (0, \epsilon)$. One can consider the limit module $\pi_0^{\mathcal{F}} = \lim_{t \rightarrow 0^+} \pi_t^{\mathcal{F}}$. This is a tempered \mathbb{H}_{k_0} -module with the same W -structure as $\pi_t^{\mathcal{F}}$. In particular, the set $\{\pi_0^{\mathcal{F}}|_W : \mathcal{F}\}$ is orthonormal in $\overline{R}_{\mathbb{Z}}(W)$, and again by the isometry r from (2.6.1), the set $\mathcal{B} = \{\pi_0^{\mathcal{F}} : \mathcal{F}\}$ is orthonormal in $\overline{R}_{\mathbb{Z}}(\mathbb{H})$. By [CKK, Theorem A], when R is of type B_n this set consists of irreducible \mathbb{H}_{k_0} -modules. Corollary 2.11 implies in particular that $\dim \overline{R}_{\mathbb{Z}}(\mathbb{H}_{k_0}) = \dim \overline{R}_{\mathbb{Z}}(\mathbb{H}_{k_t})$, and thus \mathcal{B} is a basis of $\dim \overline{R}_{\mathbb{Z}}(\mathbb{H}_{k_0})$.

5.2. $R = D_n$. We begin by recalling the definition of quasidistinguished nilpotent elements and presenting their classification when \mathfrak{g} is classical. Let e be a nilpotent element in \mathfrak{g} , and assume that e is contained in a Levi subalgebra \mathfrak{m} of \mathfrak{g} with Levi subgroup M of G . The natural map of component groups $A_M(e) \rightarrow A_G(e)$ is in fact an injection. The element e is called quasidistinguished if

$$A_G(e) \neq \bigcup_{e \in \mathfrak{m}} A_M(e), \quad (5.2.1)$$

where the union in the right hand side is over all proper Levi subalgebras containing e . Every distinguished nilpotent element is automatically quasidistinguished. The classification of nilpotent adjoint orbits when \mathfrak{g} is of classical type is based on the Jordan canonical form. The nilpotent orbits are parametrized by:

- (i) partitions of n , when $\mathfrak{g} = \mathfrak{sl}(n)$;
- (ii) partitions of $2n$ where every odd part occurs with even multiplicity, when $\mathfrak{g} = \mathfrak{sp}(2n)$;
- (iii) partitions of m where every even part occurs with even multiplicity, when $\mathfrak{g} = \mathfrak{so}(m)$, except there are two distinct nilpotent orbits in $\mathfrak{so}(2n)$ for every partition where all parts are even.

The distinguished orbits are parametrized by:

- (i) the partition (n) (principal nilpotent orbit), when $\mathfrak{g} = \mathfrak{sl}(n)$;
- (ii) partitions of $2n$ of the form (a_1, a_2, \dots, a_l) , where $0 < a_1 < a_2 < \dots < a_l$ are all even, when $\mathfrak{g} = \mathfrak{sp}(2n)$;
- (iii) partitions of m of the form (a_1, a_2, \dots, a_l) , where $0 < a_1 < a_2 < \dots < a_l$ are all odd, when $\mathfrak{g} = \mathfrak{so}(m)$.

The centralizers of nilpotent elements e and the component groups $A(e)$ are known explicitly for classical group, they were computed by Springer and Steinberg, see [Ca, pp.398-399]. A case-by-case analysis leads to the following classification of quasidistinguished orbits.

Lemma 5.2. (i) *If $\mathfrak{g} = \mathfrak{sl}(n)$, the only quasidistinguished nilpotent orbit is the principal one.*
(ii) *If $\mathfrak{g} = \mathfrak{sp}(2n)$, the quasidistinguished nilpotent orbits are labeled by partitions of $2n$ with even parts such that the multiplicity of every part is at most two.*

- (iii) If $\mathfrak{g} = \mathfrak{so}(2n+1)$, the quasidistinguished nilpotent orbits are labeled by partitions of $2n+1$ with odd parts such that the multiplicity of every part is at most two.
- (iv) If $\mathfrak{g} = \mathfrak{so}(2n)$, the quasidistinguished nilpotent orbits are labeled by partitions of $2n$ with odd parts, such that the multiplicity of every part is at most two, and if there are no parts with multiplicity one, then the number of distinct odd parts is even.

In light of Corollary 2.10, we only need to compute the elliptic norms of tempered modules $\pi_{e,\psi}$, when e is quasidistinguished, but not distinguished. By Corollary 2.11 and [R, Proposition 3.4.3]

$$\langle \pi_{e,\psi}, \pi_{e,\psi'} \rangle_{\mathbb{H}}^{\text{EP}} = \langle \pi_{e,\psi}|_W, \pi_{e,\psi'}|_W \rangle_W^{\text{ell}} = \langle \psi, \psi' \rangle_{A(e)}^{\text{ell}} \quad (5.2.2)$$

where the elliptic pairing in the right hand side of (5.2.2) is with respect to the action of $A(e)$ on the toral Lie algebra \mathfrak{s}_0 of the reductive centralizer of e in G , cf. [R, section 3.2]. Denote this latter elliptic space by $\overline{R}_{\mathbb{Z}}(A(e))$.

Lemma 5.3. *Let R be of type D_n and e be a quasidistinguished, not distinguished nilpotent element in $\mathfrak{g} = \mathfrak{so}(2n)$, parameterized by a partition τ of $2n$.*

- (a) *If $\tau = (a_1, a_1, a_2, a_2, \dots, a_l, a_l, b_1, b_2, \dots, b_{2k})$, where $k \geq 1$, $0 < a_1 < a_2 < \dots < a_l$, $0 < b_1 < b_2 < \dots < b_{2k}$, $a_i \neq b_j$, for all i, j , and all a_i and b_j are odd, then $\dim \overline{R}_{\mathbb{Z}}(A(e)) = 1$, and $\langle \text{triv}, \text{triv} \rangle_{A(e)}^{\text{ell}} = 1$.*
- (b) *If $\tau = (a_1, a_1, a_2, a_2, \dots, a_{2l}, a_{2l})$, where $0 < a_1 < a_2 < \dots < a_{2l}$, and all a_i are odd, then $\dim \overline{R}_{\mathbb{Z}}(A(e)) = 1$, and $\langle \text{triv}, \text{triv} \rangle_{A(e)}^{\text{ell}} = 2$.*

Proof. The claims follow immediately once we describe explicitly the action of $A(e)$ on \mathfrak{s}_0 .

- (a) In this case, the component group is $A(e) = (\mathbb{Z}/2\mathbb{Z})^l \times (\mathbb{Z}/2\mathbb{Z})^{2k-2}$ acting on the space \mathfrak{s}_0 of dimension l . The action is as follows: $(\mathbb{Z}/2\mathbb{Z})^{2k-2}$ acts trivially, while $(\mathbb{Z}/2\mathbb{Z})^l$ acts via

$$\bigoplus_{i=1}^l \text{triv} \boxtimes \dots \boxtimes \text{triv} \boxtimes \text{sgn}_i \boxtimes \text{triv} \boxtimes \dots \boxtimes \text{triv},$$

where sgn_i is the sgn representation on the i -th position.

- (b) In this case, the component group is $A(e) = (\mathbb{Z}/2\mathbb{Z})^{2l-1}$ acting on the space \mathfrak{s}_0 of dimension $2l$. The action is via

$$2 \bigoplus_{i=1}^{2l-1} \text{triv} \boxtimes \dots \boxtimes \text{triv} \boxtimes \text{sgn}_i \boxtimes \text{triv} \boxtimes \dots \boxtimes \text{triv}.$$

□

5.3. R of type E . It remains to discuss the cases when R is of type E_6, E_7, E_8 .

In type E_8 , there are four quasidistinguished, non-distinguished nilpotent orbits, and in all cases $A(e) \cong \mathbb{Z}/2\mathbb{Z}$ acts by the sgn representation on \mathfrak{s}_0 . This means that $\dim \overline{R}_{\mathbb{Z}}(A(e)) = 1$ and $\langle \text{triv}, \text{triv} \rangle_{A(e)}^{\text{ell}} = 1$ in all cases. The explicit relations between tempered modules are:

- (a) $D_5 + A_2$: $\pi_{D_5+A_2, \text{triv}} \oplus \pi_{D_5+A_2, \text{sgn}} = \text{ind}_{\mathbb{H}(D_5+A_2)}^{\mathbb{H}(E_8)}(\pi_{(91), \text{triv}} \boxtimes \pi_{(3), \text{triv}});$
- (b) $D_7(a_1)$: $\pi_{D_7(a_1), \text{triv}} \oplus \pi_{D_7(a_1), \text{sgn}} = \text{ind}_{\mathbb{H}(D_7)}^{\mathbb{H}(E_8)}(\pi_{(11,3), \text{triv}});$
- (c) $D_7(a_2)$: $\pi_{D_7(a_2), \text{triv}} \oplus \pi_{D_7(a_2), \text{sgn}} = \text{ind}_{\mathbb{H}(D_7)}^{\mathbb{H}(E_8)}(\pi_{(9,5), \text{triv}});$

- (d) $E_6(a_1) + A_1$: $\pi_{E_6(a_1)A_1, \text{triv}} \oplus \pi_{E_6(a_1)A_1, \text{sgn}} = \text{ind}_{\mathbb{H}(E_6+A_1)}^{\mathbb{H}(E_8)}(\pi_{E_6(a_1), \text{triv}} \boxtimes \pi_{(2), \text{triv}})$.

In type E_7 , there are two quasidistinguished, non-distinguished nilpotent orbit and in both cases $A(e) \cong \mathbb{Z}/2\mathbb{Z}$:

- (1) $E_6(a_1)$: $\pi_{E_6(a_1), \text{triv}} \oplus \pi_{E_6(a_1), \text{sgn}} = \text{ind}_{\mathbb{H}(E_6)}^{\mathbb{H}(E_7)}(\pi_{E_6(a_1), \text{triv}})$; the group $A(e)$ acts by sgn on \mathfrak{s}_0 ;
 (2) $A_4 + A_1$: $\pi_{A_4+A_1, \text{triv}} \oplus \pi_{A_4+A_1, \text{sgn}} = \text{ind}_{A_4+A_1}^{E_7}(\pi_{(5)} \boxtimes \pi_{(2)})$. Here \mathfrak{s}_0 is two dimensional, and the group $A(e) = \mathbb{Z}/2\mathbb{Z}$ acts by 2sgn on \mathfrak{s}_0 . This has the effect that $\wedge^\pm \mathfrak{s}_0 = 2(\text{triv} - \text{sgn})$, and therefore $\dim \overline{R}_{\mathbb{Z}}(A(e)) = 1$, but $\langle \text{triv}, \text{triv} \rangle_{A(e)}^{\text{ell}} = 2$.

In type E_6 , there is one quasidistinguished, non-distinguished nilpotent orbit denoted $D_4(a_1)$, whose component group is S_3 . There are three tempered modules $\pi_{D_4(a_1), \text{triv}}$, $\pi_{D_4(a_1), (21)}$, and $\pi_{D_4(a_1), \text{sgn}}$. First we have $\text{ind}_{\mathbb{H}(D_4)}^{\mathbb{H}(E_6)}(\pi_{(53), \text{triv}}) = \pi_{D_4(a_1), \text{triv}} \oplus 2\pi_{D_4(a_1), (21)} \oplus \pi_{D_4(a_1), \text{sgn}}$. Next, in D_5 , the induced module $\text{ind}_{\mathbb{H}(D_4)}^{\mathbb{H}(D_5)}(\pi_{(53), \text{triv}})$ splits into a sum of two tempered modules $\pi_{(5311), \text{triv}} \oplus \pi_{(5311), \text{sgn}}$, and we have

$$\begin{aligned} \text{ind}_{\mathbb{H}(D_5)}^{\mathbb{H}(E_6)}(\pi_{(5311), \text{triv}}) &= \pi_{D_4(a_1), \text{triv}} \oplus \pi_{D_4(a_1), (21)}, \\ \text{ind}_{\mathbb{H}(D_5)}^{\mathbb{H}(E_6)}(\pi_{(5311), \text{sgn}}) &= \pi_{D_4(a_1), (21)} \oplus \pi_{D_4(a_1), \text{sgn}}. \end{aligned} \quad (5.3.1)$$

The group $A(e) = S_3$ acts on the two dimensional \mathfrak{s}_0 via the reflection representation, therefore $\dim \overline{R}_{\mathbb{Z}}(A(e)) = 1$ and $\langle \text{triv}, \text{triv} \rangle_{A(e)}^{\text{ell}} = 1$.

This concludes the proof of Theorem 5.1.

5.4. Dirac indices. We end the section with the calculation of Dirac indices in the cases when the basis elements in $\mathcal{B}(\overline{R}_{\mathbb{Z}}(\mathbb{H}))$ have elliptic norm $\sqrt{2}$.

Example 5.4. Assume $R = E_7$ and $\delta = \pi_{A_4+A_1, \text{triv}}|_W \in \overline{R}_{\mathbb{Z}}(W)$. Then there exist genuine \widetilde{W} -representations $\widetilde{\delta}^+, \widetilde{\delta}^-, \widetilde{\delta}^- = \widetilde{\delta}^+ \otimes \text{sgn}$ such that $i(\delta) = \widetilde{\delta}^+ - \widetilde{\delta}^-$. In this case, $\widetilde{\delta}^+$ is the sum of two irreducible 64-dimensional \widetilde{W} -representations.

Proof. The first claim follows from Theorem 4.2(2) since $\langle \delta, \delta \rangle_W^{\text{ell}} = 2$. The two irreducible \widetilde{W} -representations that enter are explicitly known by [C, Table 4], where they are denoted by 64_s and 64_{ss} .

The index $i(\delta) = I(\pi_{A_4+A_1, \text{triv}})$ can also be calculated more directly as follows. Firstly, by Corollary 4.6 and (3.3.2), the only irreducible \widetilde{W} -representations $\widetilde{\sigma}$ that can contribute to $I(\pi_{A_4+A_1, \text{triv}})$ have the property that $\widetilde{\sigma}(\Omega_{\widetilde{W}}) = \langle h/2, h/2 \rangle$, where h is a middle element of a Lie triple for the nilpotent orbit $A_4 + A_1$. In [C, Table 4], it is calculated that there are only four \widetilde{W} -representations with this property: $64_s, 64_{ss}, 64_s \otimes \text{sgn}$ and $64_{ss} \otimes \text{sgn}$. Next, one needs to see which of them occur in $\pi_{A_4+A_1, \text{triv}} \otimes S^\pm$. As it is well-known, $\pi_{A_4+A_1, \text{triv}}|_W \cong H^\bullet(\mathbf{B}_e)^{\text{triv}} \otimes \text{sgn}$, where e is a nilpotent element of type $A_4 + A_1$, $H^\bullet(\mathbf{B}_e)$ is the total cohomology of the Springer fiber \mathbf{B}_e . The structure of $H^\bullet(\mathbf{B}_e)^{\text{triv}}$ is known explicitly by [BS], but we will not need this. The top degree component is the $W(E_7)$ -representation denoted by $\phi_{512,11}$ in [Ca]. Thus, Springer theory tells us that

$$\dim \text{Hom}_{W(E_7)}(\phi_{512,12}, \pi_{A_4+A_1, \text{triv}}) = 1. \quad (5.4.1)$$

(Again, in the notation of [Ca], $\phi_{512,12} = \phi_{512,11} \otimes \text{sgn}$.) A direct calculation using the software package `chevie` in the computer algebra system GAP3.4 and

the character table for $\widetilde{W}(E_7)$ from [M], reveals that

$$64_s \otimes S^+ = 64_{ss} \otimes S^+ = \phi_{512,12}. \quad (5.4.2)$$

(Notice that $\dim S^+ = 2^3$ indeed.) But this means that

$$\dim \operatorname{Hom}_{\widetilde{W}(E_7)}(64_s, \pi_{A_4+A_1, \text{triv}} \otimes S^+) = \dim \operatorname{Hom}_{\widetilde{W}(E_7)}(64_{ss}, \pi_{A_4+A_1, \text{triv}} \otimes S^+) = 1. \quad (5.4.3)$$

Since $\pi_{A_4+A_1, \text{triv}}$ is a unitary \mathbb{H} -module, [BCT, Proposition 4.9] implies that in fact

$$\ker D^+ = 64_s + 64_{ss}; \quad (5.4.4)$$

similarly, one deduces that $\ker D^- = 64_s \otimes \operatorname{sgn} + 64_{ss} \otimes \operatorname{sgn}$, and the claim follows. \square

Example 5.5. Assume $R = D_{2n}$, and $\tau = (a_1, a_1, a_2, a_2, \dots, a_{2l}, a_{2l})$ is a partition of $2n$, where $0 < a_1 < a_2 < \dots < a_{2l}$, and all a_i are odd. Then there exist genuine \widetilde{W}' -representations $\widetilde{\delta}^+, \widetilde{\delta}^-, \widetilde{\delta}^- = \operatorname{Sg}(\widetilde{\delta}^+)$ such that $i(\delta) = \widetilde{\delta}^+ - \widetilde{\delta}^-$. In this case, $\widetilde{\delta}^+$ is the sum of two irreducible \widetilde{W}' -representations.

Proof. To determine $\widetilde{\delta}_1^+$ and $\widetilde{\delta}_2^+$ explicitly, we rely on the classification of genuine $\widetilde{W}(D_m)$ -modules [Re], and the Dirac cohomology calculations in [C]. Recall that a complete set of inequivalent genuine irreducible $\widetilde{W}(B_m)$ -modules is given by $\{(\sigma \times \emptyset) \otimes \mathcal{S} : \sigma \text{ partition of } m\}$, where $\mathcal{S} \in \{S^+, S^-\}$, when m is odd, and $\mathcal{S} = S$, when m is even. The restriction to $\widetilde{W}(D_m)$ yields the following complete sets of inequivalent irreducible modules:

- (i) $\{(\sigma \times \emptyset) \otimes S^+ : \sigma \text{ partition of } m\}$, when m is odd. In this case, $(\sigma \times \emptyset) \otimes S^- = (\sigma^t \times \emptyset) \otimes S^+$ as $\widetilde{W}(D_m)$ -representations.
- (ii) $\{(\sigma \times \emptyset) \otimes S : \sigma \text{ partition of } m, \sigma^t \neq \sigma\} \cup \{\widetilde{\sigma}_1, \widetilde{\sigma}_2 : \sigma \text{ partition of } m, \sigma^t = \sigma\}$, when m is even, where $(\sigma \times \emptyset) \otimes S|_{W(D_m)} = \widetilde{\sigma}_1 \oplus \widetilde{\sigma}_2$, when $\sigma^t = \sigma$. In this case, every irreducible $\widetilde{W}(D_m)$ -representation is self dual under tensoring with sgn .

Returning to our example τ in D_{2n} , consider the strings $(\frac{a_i-1}{2}, \frac{a_i-3}{2}, \dots, -\frac{a_i-1}{2})$, $1 \leq i \leq 2l$. (These strings give the standard coordinates of $h/2$, where h is the middle element of a Lie triple attached to τ .) There is one way to form a partition σ of n such that these strings form the hooks of a left-justified decreasing tableau with shape σ and content $i-j$ for the (i, j) -box. For example, when $\tau = (3, 3, 5, 5)$ in D_8 , the partition σ is $(3, 3, 2)$, see Figure 1.

0	1	2
-1	0	1
-2	-1	

FIGURE 1. Nilpotent $\tau = (3, 3, 5, 5)$ and partition $\sigma = (3, 3, 2)$.

Notice that such σ always has the property that $\sigma^t = \sigma$. By [C], it follows that $\widetilde{\delta}^+ = \widetilde{\sigma}_1^+ \oplus \widetilde{\sigma}_2^+$ as $\widetilde{W}(D_{2n})'$ -representations. \square

5.5. Integrality properties of Ind_D . Recall the sublattice $\mathcal{Y} = \mathfrak{r}(\overline{R}_{\mathbb{Z}}(\mathbb{H})) \subset \overline{R}_{\mathbb{Z}}(W)$ defined in (4.4.6). As in the proof of Corollary 4.13, the restriction

$$\text{Ind}_D|_{\mathcal{Y}} : \mathcal{Y} \rightarrow \overline{R}_{\mathbb{Z}}(\mathbb{H})$$

is an isometric isomorphism and an inverse of $\mathfrak{r} : \overline{R}_{\mathbb{Z}}(\mathbb{H}) \rightarrow \mathcal{Y}$. A natural question is when

$$\mathcal{Y} = \overline{R}_{\mathbb{Z}}(W),$$

or equivalently when Ind_D gives an isomorphism $\text{Ind}_D : \overline{R}_{\mathbb{Z}}(W) \rightarrow \overline{R}_{\mathbb{Z}}(\mathbb{H})$.

Proposition 5.6. *Suppose R is irreducible. If one of the following conditions is satisfied:*

- (1) *the parameter function k is constant;*
- (2) *R is non-simply laced and the parameter k is generic;*
- (3) *R is B_n or G_2 ,*

then $\mathcal{Y} = \overline{R}_{\mathbb{Z}}(W)$.

Proof. (1) Assume without loss of generality that $k \equiv 1$. Then consider the restriction of the orthogonal basis $\mathcal{B}(\overline{R}_{\mathbb{Z}}(\mathbb{H}))$ from Theorem 5.1 to $\overline{R}_{\mathbb{Z}}(W)$. Then $\mathfrak{r}(\mathcal{B}(\overline{R}_{\mathbb{Z}}(\mathbb{H})))$ is an orthogonal set and an \mathbb{R} -basis for $\overline{R}_{\mathbb{Z}}(W)$. To see that in fact it is a \mathbb{Z} -basis, it is sufficient to recall that the geometric realization of tempered modules of the Hecke algebra ([KL, L2]) implies that the matrix of restrictions to W of the set of irreducible tempered modules with real central character is upper uni-triangular in the natural ordering given by the closure ordering of nilpotent orbits, see [BM, section 4].

In cases (2) and (3), the basis of $\overline{R}_{\mathbb{Z}}(\mathbb{H})$ constructed in Theorem 5.1 is in fact an orthonormal basis given by discrete series for generic parameters and limits of discrete series for special parameters. Thus the restriction to $\overline{R}_{\mathbb{Z}}(W)$ forms an orthonormal \mathbb{Z} -basis as well. □

Proposition 5.6(1,2), together with Corollary 4.13(2) and Theorem 4.18, allow us to improve the result in Corollary 4.13(4).

Corollary 5.7. *If $\delta \in \overline{R}_{\mathbb{Z}}(W)$ is a rational multiple of a pure element in $\overline{R}_{\mathbb{Z}}(W)$, then $\text{Ind}_D(\delta)$ is supported at a single central character.*

Proof. We only need to prove the claim when R is non-simply laced and the parameter function k is specialized to a non-generic point k_0 , since otherwise by Proposition 5.6(1,2) the claim is equivalent with Corollary 4.13(4). Assume this is the case and that X is an irreducible \mathbb{H} -module such that $\langle \text{Ind}_D(\delta), X \rangle_{\mathbb{H}}^{\text{EP}} \neq 0$. By Theorem 4.11(2), this means that $\langle i(\delta), I(X) \rangle_{\widetilde{W}'} \neq 0$, and therefore, by Corollary 4.6 the central character of X equals $\chi^{\tilde{\delta}}$ for some irreducible component $\tilde{\delta}$ of $i(\delta)$.

By Theorem 3.2, we know that $\chi^{\tilde{\delta}}$ depends polynomially on the parameter k . Moreover, at generic k , δ is a pure element in \mathcal{Y} (since $\mathcal{Y} = \overline{R}_{\mathbb{Z}}(W)$) and so, by Corollary 4.13(4), $\chi^{\tilde{\delta}}$ are equal to each other as functions of k for all constituents $\tilde{\delta}$ of $i(\delta)$. (Here, we implicitly use that Corollary 4.13(2) gives $I(\text{Ind}_D(\delta)) = i(\delta)$, independently of k .) But then the $\chi^{\tilde{\delta}}$ are equal to each other at k_0 as well. □

We end the section with a calculation of $\text{Ind}_D(\delta)$, for the basis elements $\delta \in \overline{R}_{\mathbb{Z}}(W)$ which are pure, but no units. These are the cases appearing in E_7 and D_{2n} in Theorem 5.1.

Example 5.8. Retain the notation from Example 5.4. Let $\delta = \pi_{A_4+A_1, \text{triv}}|_W$. Then

$$\text{Ind}_D(\delta) = 2\Phi([\mathbb{H} \otimes_W (\phi_{512,12} - \phi_{512,11})]).$$

We remark that $\phi_{512,12}$ and $\phi_{512,11}$ are the irreducible Springer representations attached to $(A_4 + A_1, \text{triv})$ and $(A_4 + A_1, \text{sgn})$, respectively.

Proof. From Example 5.4, we see that $\tilde{\delta}^+ = 64_s + 64_{ss}$, and thus $\text{Ind}_D(\delta) = \Phi([\mathbb{H} \otimes_W (64_s + 64_{ss}) \otimes (S^+ - S^-)]) = 2\Phi([\mathbb{H} \otimes_W (\phi_{512,12} - \phi_{512,11})])$, by (5.4.2). \square

Example 5.9. Retain the notation from Example 5.5. Let $\delta = \pi_{(1133), \text{triv}}|_W$ in $\overline{R}_{\mathbb{Z}}(W(D_4))$. Then

$$\text{Ind}_D(\delta) = 2\Phi([\mathbb{H} \otimes_W (11 \times 2 + 22 \times 0 - 12 \times 1)]),$$

where the notation for irreducible $W(D_4)$ -representations is as in [Ca]. We remark that 12×1 and 11×2 are the irreducible Springer representations attached to $((1133), \text{triv})$ and $((1133), \text{sgn})$, respectively.

Proof. From Example 5.5, we see that

$$(22 \times 0) \otimes S = \tilde{\sigma}_1 + \tilde{\sigma}_2, \text{ as } \widetilde{W}(D_4)\text{-representations,}$$

where S is the unique spin module for D_4 (of dimension 4) $\tilde{\sigma}_1, \tilde{\sigma}_2$ are two sign self-dual irreducible $\widetilde{W}(D_4)$ -representations of dimension 4. Then $\tilde{\sigma}_i = \tilde{\sigma}_i^+ + \tilde{\sigma}_i^-$ as $\widetilde{W}(D_4)'$ -representations, and the index of $\pi_{(1133), \text{triv}}$ is the virtual $\widetilde{W}(D_4)'$ -representation:

$$\begin{aligned} i(\delta) &= I(\pi_{(1133), \text{triv}}) = (\tilde{\sigma}_1^+ + \tilde{\sigma}_2^+) - (\tilde{\sigma}_1^- + \tilde{\sigma}_2^-), \text{ while} \\ I(\pi_{(1133), \text{sgn}}) &= (\tilde{\sigma}_1^- + \tilde{\sigma}_2^-) - (\tilde{\sigma}_1^+ + \tilde{\sigma}_2^+). \end{aligned} \quad (5.5.1)$$

Set $\tilde{\delta}^+ = \tilde{\sigma}_1^+ + \tilde{\sigma}_2^+$ and similarly define $\tilde{\delta}^-$. By definition,

$$\text{Ind}_D(\delta) = \Phi(\mathbb{H} \otimes_{W(D_4)'} (\tilde{\delta}^+)^* \otimes (S^+ - S^-)) = \Phi(\mathbb{H} \otimes_{W(D_4)} \otimes \text{Ind}_{W(D_4)'}^{W(D_4)} (\tilde{\delta}^- \otimes (S^+ - S^-))),$$

where S^{\pm} are the two irreducible two dimensional components of $S|_{\widetilde{W}(D_4)'}$. We need to compute $\tilde{\sigma}_i^- \otimes S^{\pm}$; in fact it is sufficient to compute their induction to $W(D_4)$. Notice that all $\tilde{\sigma}_i^- \otimes S^{\pm}$ occur as components of the restriction to $W(D_4)'$ of

$$(22 \times 0) \otimes S \otimes S = (22 \times 0) \otimes \wedge^{\bullet} V = 2(12 \times 1 + 11 \times 2 + 22 \times 0).$$

Each of 12×1 , 11×2 , 22×0 is sign self-dual, hence they break into a sum of two irreducible equidimensional $W(D_4)'$ -representations of dimensions 4, 3, and 1, respectively. This means that every $\text{Ind}_{W(D_4)'}^{W(D_4)} (\tilde{\sigma}_i^{\pm} \otimes S^{\pm})$ equals either

$$12 \times 1, \text{ or } 11 \times 2 + 22 \times 0. \quad (5.5.2)$$

Using [CT, Lemma 3.8] and (5.5.1), we see that $\tilde{\sigma}_1^- + \tilde{\sigma}_2^-$ is contained in $(12 \times 1) \otimes S^+$ and in $(11 \times 2) \otimes S^+$. Combining this with (5.5.2), it follows that

$$\text{Ind}_{W(D_4)'}^{W(D_4)} (\tilde{\sigma}_i^- \otimes S^-) = 12 \times 1 \text{ and } \text{Ind}_{W(D_4)'}^{W(D_4)} (\tilde{\sigma}_i^- \otimes S^+) = 11 \times 2 + 22 \times 0, \quad i = 1, 2,$$

and this proves the claim. \square

6. DIRAC INDUCTION (ANALYTIC VERSION)

6.1. An analytic model. Let $C^\omega(V)$ be the set of complex analytic functions on V . Let M be a finite dimensional (unitary) W -module. Following [HO, EOS] we define a left module action of \mathbb{H} on $C^\omega(V) \otimes_{\mathbb{C}} M$. Moreover, we define an inner product on $C^\omega(V) \otimes M$ which makes this (actually the subset of “finite” vectors) into a unitary \mathbb{H} -module, with respect to the natural $*$ -operation on \mathbb{H} from section 3.2.

We begin by defining certain operators on $C^\omega(V) \otimes M$.

Definition 6.1. For every $v \in V$, $f \in C^\omega(V)$, let ∂_v denote the directional derivative of f in the direction of v . Let $Q(v) : C^\omega(V) \otimes M \rightarrow C^\omega(V) \otimes M$ be the operator

$$Q(v)(f \otimes m) = \partial_v f \otimes m. \quad (6.1.1)$$

The Weyl group W acts naturally on $C^\omega(V)$ via the left regular action $(wf)(\xi) = f(w^{-1}\xi)$. For every root $\alpha \in R$, define the integral operator $I(\alpha) : C^\omega(V) \rightarrow C^\omega(V)$,

$$I(\alpha)f(\xi) = \int_0^{(\xi, \alpha^\vee)} f(\xi - t\alpha) dt, \quad \xi \in V. \quad (6.1.2)$$

For every $\alpha \in F$, let $Q(s_\alpha) : C^\omega(V) \otimes M \rightarrow C^\omega(V) \otimes M$ be the operator

$$Q(s_\alpha)(f \otimes m) = s_\alpha f \otimes s_\alpha m + k_\alpha I(\alpha)f \otimes m. \quad (6.1.3)$$

Theorem 6.2 ([EOS, Theorem 4.11]¹). *The assignment $v \rightarrow Q(v)$, $s_\alpha \rightarrow Q(s_\alpha)$ extends to an action (the “integral-reflection” representation) of \mathbb{H} on $C^\omega(V) \otimes M$.*

The explanation is as follows ([EOS, section 4.3]). Start with the induced module $\mathbb{H} \otimes_{\mathbb{C}[W]} M^*$ with the action of \mathbb{H} by left multiplication. As \mathbb{C} -vector spaces, $\mathbb{H} \otimes_{\mathbb{C}[W]} M^*$ is isomorphic with $S(V_{\mathbb{C}}) \otimes_{\mathbb{C}} M^*$. One traces the action under this identification. For $\alpha \in F$, $v \in V$, $m' \in M^*$:

$$\begin{aligned} s_\alpha(p \otimes m') &= (s_\alpha \cdot p) \otimes m' = (s_\alpha(p)s_\alpha + k_\alpha \Delta_\alpha(p)) \otimes m' \\ &= s_\alpha(p) \otimes s_\alpha m' + k_\alpha \Delta_\alpha(p) \otimes m'; \\ v(p \otimes m') &= vp \otimes m', \end{aligned}$$

where Δ_α is the difference operator from (2.3.1).

Now consider the dual $(S(V_{\mathbb{C}}) \otimes M^*)^* \supset C^\omega(V) \otimes M$. The previous action of \mathbb{H} on $S(V_{\mathbb{C}}) \otimes M^*$ defines an action of \mathbb{H} on the dual $(S(V_{\mathbb{C}}) \otimes M^*)^*$ by means of the anti-automorphism \star defined on the generators of \mathbb{H} by $w^\star = w^{-1}$ and $\xi^\star = \xi$. To get to the integral-reflection action Q from Theorem 6.2, one uses the fact that under the natural pairing

$$(\cdot, \cdot) : S(V_{\mathbb{C}}) \otimes C^\omega(V) \rightarrow \mathbb{C}, (p, f) = (p(\partial)f)(0),$$

the operator Δ_α is adjoint to the operator $I(\alpha)$.

¹In fact, [EOS] defines an action of the trigonometric Cherednik algebra at critical level (which contains \mathbb{H} as a subalgebra).

6.2. Unitary structure. We generalize the inner product from [HO, (2.6)]. To begin, for every Weyl chamber C , define an inner product $(\ , \)_C$ on $C^\omega(V) \otimes M$ by extending linearly

$$(\psi_1 \otimes m_1, \psi_2 \otimes m_2)_C = (\psi_1, \psi_2)_C (m_1, m_2)_M, \quad \psi_1, \psi_2 \in C^\omega(V), m_1, m_2 \in M, \quad (6.2.1)$$

where $(\psi_1, \psi_2)_C = \int_C \psi_1(\eta) \overline{\psi_2(\eta)} d\eta$, and $(\ , \)_M$ is a fixed W -invariant inner product on M .

Then, for every $f, g \in C^\omega(V) \otimes M$, set

$$(f, g)_{k,C} = \sum_{w \in W} (Q(w)f, Q(w)g)_C. \quad (6.2.2)$$

The notation is meant to emphasize that this inner product depends on the multiplicity function k (since Q does) and on the choice of chamber C . Let C_+ be the fundamental Weyl chamber (corresponding to F).

Theorem 6.3. *The inner product $(\ , \)_{k,C_+}$ is $*$ -invariant for \mathbb{H} .*

Proof. The proof is an adaptation of the proof of [HO, Theorem 2.4].

The invariance with respect to w is clear since the inner product averages over W . Fix $\xi \in V$. A formal argument, using only the definition of $*$ and the relations in the Hecke algebra shows that

$$(\partial(\xi)f, g)_{k,C} - (f, \partial(\xi)^*g)_{k,C} = \sum_w (\partial(w\xi)Q(w)f, Q(w)g)_C + \sum_w (Q(w)f, \partial(w\xi)Q(w)g)_C.$$

Write $Q(w)f = \sum_j \phi_{w,j} \otimes m_{w,j}$ and $Q(w)g = \sum_i \psi_{w,i} \otimes n_{w,i}$, where $\phi_{w,j}, \psi_{w,i} \in C^\omega(V)$ and $m_{w,j}, n_{w,i} \in M$. Set $h_{w,i,j}(\eta) = \phi_{w,j}(\eta) \overline{\psi_{w,i}(\eta)}$. Then:

$$\begin{aligned} & (\partial(w\xi)Q(w)f, Q(w)g)_C + (Q(w)f, \partial(w\xi)Q(w)g)_C \\ &= \sum_{i,j} \int_C (\partial(w\xi)\phi_{w,j}(\eta)) \overline{\psi_{w,i}(\eta)} d\eta (m_{w,j}, n_{w,i})_M \\ &+ \sum_{i,j} \phi_{w,j}(\eta) \overline{(\partial(w\xi)\psi_{w,i}(\eta))} d\eta (m_{w,j}, n_{w,i})_M \\ &= \sum_{i,j} \int_C \partial(w\xi)h_{w,i,j}(\eta) d\eta (m_{w,j}, n_{w,i})_M \\ &= \sum_{i,j} \int_{\partial C} h_{w,i,j}(\eta) (w\xi, \nu) d\sigma(\eta) (m_{w,j}, n_{w,i})_M, \end{aligned}$$

by Stokes' theorem, where ν is the outer normal vector. Now assume $C = C_+$. Since the boundary of C_+ is formed of the intersections with the root hyperplanes H_α , $\alpha \in F$, one finds that this equals

$$\sum_{i,j} \sum_{\alpha \in F} \int_{H_\alpha \cap C_+} h_{w,i,j}(\eta) (w\xi, \frac{\alpha^\vee}{|\alpha^\vee|}) d\sigma_\alpha(\eta) (m_{w,j}, n_{w,i}).$$

Summing over w and changing the order of summation, one gets:

$$\begin{aligned} \sum_{\alpha \in F} \int_{H_\alpha \cap C_+} & \left(\sum_{w^{-1}\alpha > 0} \sum_{i,j} h_{w,i,j}(\eta)(w\xi, \frac{\alpha^\vee}{|\alpha^\vee|})(m_{w,j}, n_{w,i}) + \right. \\ & \left. \sum_{w^{-1}\alpha < 0} \sum_{i,j} h_{w,i,j}(\eta)(w\xi, \frac{\alpha^\vee}{|\alpha^\vee|})(m_{w,j}, n_{w,i}) \right) d\sigma_\alpha(\eta). \end{aligned}$$

We wish to show that this quantity is zero. We make the substitution $w' = s_\alpha w$ in the second sum since then $(w\xi, \frac{\alpha^\vee}{|\alpha^\vee|}) = -(w'\xi, \frac{\alpha^\vee}{|\alpha^\vee|})$ and $(w')^{-1}\alpha > 0$ if $w^{-1}\alpha < 0$. But it remains to verify how $h_{w,i,j}, m_{w,j}, n_{w,i}$ are related to $h_{w',i',j'}, m_{w',j'}, n_{w',i'}$ on the hyperplane H_α . For this, notice that $I(\alpha)h = 0$ and $s_\alpha \cdot h = h$, on H_α for all $h \in C^\omega(V)$. Therefore, on H_α :

$$Q(w')f = Q(s_\alpha) \sum_j \phi_{w,j} \otimes m_{w,j} = \sum_j s_\alpha \phi_{w,j} \otimes s_\alpha m_{w,j} = \sum_j \phi_{w,j} \otimes s_\alpha m_{w,j}, \quad (6.2.3)$$

and similarly for $Q(w')g$. This implies that on H_α , we have $h_{w',i,j} = h_{w,i,j}$, $m_{w',j} = s_\alpha m_{w,j}$, $n_{w',i} = s_\alpha n_{w,i}$ (implicit here is that the sets of indices (i,j) for w and w' are the same). Now the claim follows by the W -invariance of the product $(\cdot, \cdot)_M$. \square

Remark 6.4. A completely similar argument shows that the inner product $(\cdot, \cdot)_{k,C_-}$ is also invariant, where $C_- = w_0(C_+)$ is the negative Weyl chamber.

Define

$$\mathcal{X}_\omega(M) = \{f \in C^\omega(V) \otimes M : (\partial(p)f, \partial(p)f)_{k,C_+} < \infty, \text{ for all } p \in S(V_\mathbb{C})\}. \quad (6.2.4)$$

Using the W -invariance of $(\cdot, \cdot)_{k,C_+}$, it is easy to see that the action of \mathbb{H} preserves $\mathcal{X}_\omega(M)$. Thus we have:

Corollary 6.5. *The \mathbb{H} -module $\mathcal{X}_\omega(M)$ is $*$ -pre-unitary.*

6.3. Global Dirac operators. We wish to define Dirac operators on the spaces $\mathcal{X}_\omega(M)$. For this we need to trace again through the action of \mathbb{H} and the chain of identifications after Theorems 6.2.

Let S be a spin module for the Clifford algebra $C(V)$ and let E be a genuine \widetilde{W} -module. Then $E \otimes S$ is a W -representation, so we have the \mathbb{H} -module $\mathcal{X}_\omega(E \otimes S)$.

If \mathcal{B} is an orthonormal basis of V , define $D \in \text{End}_{\mathbb{H}}(\mathbb{H} \otimes_W (E \otimes S))$ via

$$D(h \otimes x \otimes y) = \sum_{\xi \in \mathcal{B}} h \widetilde{\xi} \otimes x \otimes \xi y, \quad h \in \mathbb{H}, x \in E, y \in S. \quad (6.3.1)$$

The definition does not depend on the choice of basis \mathcal{B} , and moreover D is well-defined:

$$\begin{aligned} D(hw \otimes w^{-1}(x \otimes y)) &= \sum_{\xi \in \mathcal{B}} hw \widetilde{\xi} \otimes \widetilde{w}^{-1}x \otimes \xi \widetilde{w}^{-1}y \quad (\text{for some pullback } \widetilde{w} \text{ of } w \text{ in } \widetilde{W}) \\ &= \sum_{\xi \in \mathcal{B}} h \widetilde{w(\xi)} w \otimes \widetilde{w}^{-1}x \otimes \widetilde{w}^{-1}w(\xi)y \quad (\text{where } \widetilde{w(\xi)} = w \cdot \widetilde{\xi} \cdot w^{-1}) \\ &= \sum_{\xi \in \mathcal{B}} h \widetilde{w(\xi)} w \otimes w^{-1}(x \otimes w(\xi)y) = \sum_{\xi \in w(\mathcal{B})} h \widetilde{\xi} \otimes (x \otimes \xi y) = D(h \otimes x \otimes y), \end{aligned}$$

since $w(\mathcal{B})$ is also an orthonormal basis of V .

Clearly D commutes with the module action of \mathbb{H} , since the \mathbb{H} -action is by left multiplication.

In the identification $\mathbb{H} \otimes_W (E \otimes S) \cong S(V_{\mathbb{C}}) \otimes (E \otimes S)$, D acts as follows:

$$\begin{aligned} D(p \otimes (x \otimes y)) &= \sum_{\xi \in \mathcal{B}} p \tilde{\xi} \otimes (x \otimes \xi y) = \sum_{\xi \in \mathcal{B}} (p \xi \otimes (x \otimes \xi y) - p T_{\xi} \otimes (x \otimes \xi y)) \\ &= \sum_{\xi \in \mathcal{B}} \xi p \otimes (x \otimes \xi y) - \sum_{\xi \in \mathcal{B}} p \otimes T_{\xi}(x \otimes \xi y), \end{aligned}$$

where T_{ξ} were defined in (3.2.2). The formula for D^2 can be computed analogously with the one in the local case (3.2.6).

Proposition 6.6. *As operators on $\mathbb{H} \otimes_W (E \otimes S)$, we have*

$$D^2 = -\Omega \otimes 1 \otimes 1 + 1 \otimes \Omega_{\widetilde{W}} \otimes 1. \quad (6.3.2)$$

Proof. The proof is a completely analogous calculation to that in the proof of [BCT, Theorem 3.5]. For simplicity, denote

$$R_{\circ}^2 = \{(\alpha, \beta) \in R \times R : \alpha, \beta > 0, \alpha \neq \beta, s_{\alpha}(\beta) < 0\}. \quad (6.3.3)$$

Let $\{\xi_i\}$ be an orthonormal basis of V and let $p \otimes (x \otimes y)$ be an element of $\mathbb{H} \otimes (E \otimes S)$. Then we have

$$\begin{aligned} D^2(p \otimes (x \otimes y)) &= \sum_{i,j} p \tilde{\xi}_i \tilde{\xi}_j \otimes (x \otimes \xi_j \xi_i y) \\ &= \sum_i p \tilde{\xi}_i^2 \otimes x \otimes \xi_i^2 y + \sum_{i < j} p [\tilde{\xi}_i, \tilde{\xi}_j] \otimes (x \otimes \xi_j \xi_i y) \end{aligned}$$

Now, we use the identities

$$\sum_i \tilde{\xi}_i^2 = \Omega - \frac{1}{4} \sum_{\alpha > 0} \langle \alpha^{\vee}, \alpha^{\vee} \rangle - \frac{1}{4} \sum_{(\alpha, \beta) \in R_0^2} k_{\alpha} k_{\beta} \langle \alpha^{\vee}, \beta^{\vee} \rangle s_{\alpha} s_{\beta}, \quad (\text{see [BCT, Theorem 2.11]}) \quad (6.3.4)$$

and

$$[\tilde{\xi}_i, \tilde{\xi}_j] = [T_{\xi_j}, T_{\xi_i}] = \frac{1}{4} \sum_{(\alpha, \beta) \in R_0^2} k_{\alpha} k_{\beta} ((\alpha^{\vee}, \xi_j)(\beta^{\vee}, \xi_i) - (\beta^{\vee}, \xi_j)(\alpha^{\vee}, \xi_i)) s_{\alpha} s_{\beta}, \quad (6.3.5)$$

(see for example [BCT, Lemma 2.9]). It follows that:

$$\begin{aligned} D^2(p \otimes (x \otimes y)) &= -\Omega p \otimes (x \otimes y) + \frac{1}{4} \sum_{\alpha > 0} \langle \alpha^{\vee}, \alpha^{\vee} \rangle p \otimes (x \otimes y) \\ &\quad + \frac{1}{4} \sum_{(\alpha, \beta) \in R_0^2} k_{\alpha} k_{\beta} \langle \alpha^{\vee}, \beta^{\vee} \rangle p \otimes s_{\alpha} s_{\beta} (x \otimes y) \\ &\quad + \frac{1}{4} \sum_{i < j} \sum_{(\alpha, \beta) \in R_0^2} k_{\alpha} k_{\beta} p \otimes ((\alpha^{\vee}, \xi_j)(\beta^{\vee}, \xi_i) - (\beta^{\vee}, \xi_j)(\alpha^{\vee}, \xi_i)) s_{\alpha} s_{\beta} (x \otimes \xi_j \xi_i y) \\ &= -\Omega p \otimes (x \otimes y) + \frac{1}{4} \sum_{\alpha > 0} \langle \alpha^{\vee}, \alpha^{\vee} \rangle p \otimes (x \otimes y) + \frac{1}{4} \sum_{(\alpha, \beta) \in R_0^2} k_{\alpha} k_{\beta} \langle \alpha^{\vee}, \beta^{\vee} \rangle p \otimes s_{\alpha} s_{\beta} (x \otimes y) \\ &\quad - \frac{1}{4} \sum_{i \neq j} \sum_{(\alpha, \beta) \in R_0^2} k_{\alpha} k_{\beta} p \otimes (\beta^{\vee}, \xi_j)(\alpha^{\vee}, \xi_i) s_{\alpha} s_{\beta} (x \otimes \xi_j \xi_i y), \end{aligned}$$

where we used again that $\xi_i \xi_j = -\xi_j \xi_i$ for $i \neq j$. Now changing the order of summation, the last double sum becomes

$$\frac{1}{4} \sum_{(\alpha, \beta) \in R_0^2} k_\alpha k_\beta p \otimes s_\alpha s_\beta (x \otimes (\beta^\vee \alpha^\vee + \langle \alpha^\vee, \beta^\vee \rangle) y),$$

where we identify α^\vee, β^\vee , via the inner product $\langle \cdot, \cdot \rangle$, as elements of $C(V)$. Finally, making the necessary cancellations, we arrive at the desired formula. \square

Dualizing and making the identifications, we arrive at the following definition.

Definition 6.7. Let E be a genuine \widetilde{W} -module. The Dirac operator $D_E \in \text{End}_{\mathbb{H}}(C^\omega(V) \otimes (E \otimes S))$ is given by

$$D_E(f \otimes x \otimes y) = \sum_{\xi \in \mathcal{B}} \partial(\xi) f \otimes x \otimes \xi y - \sum_{\xi \in \mathcal{B}} f \otimes T_\xi(x \otimes \xi y), \quad (6.3.6)$$

where $T_\xi = \frac{1}{2} \sum_{\beta \in R^+} k_\beta(\xi, \beta^\vee) s_\beta$. Clearly, D_E preserves the finite vectors in $C^\omega(V) \otimes (E \otimes S)$, and thus it defines an operator $D_E \in \text{End}_{\mathbb{H}}(\mathcal{X}_\omega(E \otimes S))$.

Example 6.8. We consider the Hecke algebra \mathbb{H} for $sl(2)$. Here $V = \mathbb{R}\alpha$, $W = \mathbb{Z}/2\mathbb{Z}$, and \mathbb{H} is generated by s and $\xi \in V$ subject to

$$s \cdot \xi + \xi \cdot s = \langle \xi, \alpha^\vee \rangle.$$

We assume that the inner product on V is normalized so that the length on α^\vee is $\sqrt{2}$.

If we make the identification $\xi \rightarrow \xi\alpha$, $\xi \in \mathbb{R}$, we may regard the functions f as $f : \mathbb{R} \rightarrow \mathbb{R}$, and the action of the Hecke algebra on $C^\omega(\mathbb{R}) \otimes M$ is:

$$Q(s)(f(\xi) \otimes m) = f(-\xi) \otimes (s \cdot m) + \left(\int_{-\xi}^{\xi} f(t) dt \right) \otimes m;$$

$$Q(\xi)f = \frac{df}{d\xi}.$$

The cover \widetilde{W} is isomorphic to $\mathbb{Z}/4\mathbb{Z}$ and there are two genuine \widetilde{W} -types (both spin modules), χ_+ and χ_- given by multiplication by i and $-i$ respectively. Fix $S = \chi_+$ and take the basis $\mathcal{B} = \{\frac{1}{\sqrt{2}}\alpha^\vee\}$.

- (1) $E = \chi_+$. Note that $E \otimes S$ is the **sgn** W -representation. We have $D_+(f \otimes x \otimes y) = i(\frac{df}{d\xi} \otimes x \otimes y + f \otimes x \otimes y)$. Then $\ker D_+ = \mathbb{R}e^{-\xi} \otimes \mathbf{sgn}$, and one checks that this is the Steinberg module. It is unitary with respect to the inner product $(\cdot, \cdot)_{C_+}$.
- (2) $E = \chi_-$. Note that $E \otimes S$ is the **triv** W -representation. We have $D_-(f \otimes x \otimes y) = i(\frac{df}{d\xi} \otimes x \otimes y - f \otimes x \otimes y)$. Then $\ker D_- = \mathbb{R}e^\xi \otimes \mathbf{triv}$, and one checks that this is the trivial module. It is unitary with respect to the inner product $(\cdot, \cdot)_{C_-}$.

Definition 6.9. Assume that $\dim V$ is odd, and let S^+, S^- be the two spin modules of $C(V)$. Define $D_E^\pm : \mathcal{X}_\omega(E \otimes S^\pm) \rightarrow \mathcal{X}_\omega(E \otimes S^\mp)$ by composing the Dirac operators $D_E^\pm \in \text{End}_{\mathbb{H}}(\mathcal{X}(E \otimes S^\pm))$ from Definition 6.7 with the vector space isomorphism $S^+ \rightarrow S^-$, as in the local case in section 4.1.

Now assume that $\dim V$ is even. Let S^+, S^- be the two spin modules of $C(V)_{\text{even}}$ and let E be a genuine representation of \widetilde{W}' . Then $E \otimes S^\pm$ are W' -representations, so we can consider the \mathbb{H} -modules

$$\mathcal{X}'_\omega(E \otimes S^\pm) := \mathcal{X}_\omega(\mathbb{C}[W] \otimes_{W'} S^\pm). \quad (6.3.7)$$

Define Dirac operators

$$D^\pm : \mathbb{H} \otimes_{W'} (E \otimes S^\pm) \rightarrow \mathbb{H} \otimes_{W'} (E \otimes S^\mp)$$

by

$$D^\pm(h \otimes x \otimes y) = \sum_{\xi \in \mathcal{B}} h \tilde{\xi} \otimes x \otimes \xi y, \quad h \in \mathbb{H}, x \in E, y \in S.$$

Again, D^\pm are well-defined, independent on the choice of basis \mathcal{B} , and commute with the left action of \mathbb{H} . The formula for $D^\pm D^\mp$ is the same as in the formula (6.3.2):

$$D^\pm D^\mp = -\Omega \otimes 1 \otimes 1 + 1 \otimes \Omega_{\widetilde{W}} \otimes 1, \quad (6.3.8)$$

because in that calculation, the elements of W that occur are actually in W' , so they can still be moved across the tensor product.

In the identification

$$\mathbb{H} \otimes_{W'} (E \otimes S^\pm) \cong \mathbb{H} \otimes_W (\mathbb{C}[W] \otimes_{W'} (E \otimes S^\pm)) \cong S(V_{\mathbb{C}}) \otimes (\mathbb{C}[W] \otimes_{W'} (E \otimes S^\pm)), \quad (6.3.9)$$

the action of D^\pm is as follows:

$$D^\pm(p \otimes (w \otimes (x \otimes y))) = \sum_{\xi \in \mathcal{B}} \xi p \otimes (w \otimes (x \otimes \xi y)) - \sum_{\xi \in \mathcal{B}} p \otimes (T_\xi w \otimes (x \otimes \xi y)).$$

Dualizing, we obtain the following definition.

Definition 6.10. Assume that $\dim V$ is even. Let E be a genuine \widetilde{W}' -module. The Dirac operators $D_E^\pm : \mathcal{X}'_\omega(E \otimes S^\pm) \rightarrow \mathcal{X}'_\omega(E \otimes S^\mp)$ are given by

$$D_E^\pm(f \otimes (w \otimes (x \otimes y))) = \sum_{\xi \in \mathcal{B}} \partial(\xi) f \otimes (w \otimes (x \otimes \xi y)) - \sum_{\xi \in \mathcal{B}} f \otimes (T_\xi w \otimes (x \otimes \xi y)). \quad (6.3.10)$$

6.4. Global Dirac index. With these definitions at hand, we can define the global Dirac index. For uniformity of notation, set $\mathcal{X}'_\omega(M) = \mathcal{X}_\omega(M)$, when $\dim V$ is odd as well, for every $W' = W$ -module M . For every genuine \widetilde{W}' -representation E , we have defined the Dirac operators

$$D_E^\pm : \mathcal{X}'_\omega(E \otimes S^\pm) \rightarrow \mathcal{X}'_\omega(E \otimes S^\mp), \quad (6.4.1)$$

which commute with the action of \mathbb{H} . As in the case of local Dirac operators, it is easy to see that D_E^+ is adjoint to D_E^- with respect to the unitary structure of $\mathcal{X}'_\omega(E \otimes S^\pm)$.

If $\lambda \in W \setminus V_{\mathbb{C}}^\vee$, define

$$\mathcal{X}'_\omega(M)_\lambda = \{f \in \mathcal{X}'_\omega(M) : \partial(p)f = p(\lambda)f, \text{ for all } p \in S(V_{\mathbb{C}})^W\}. \quad (6.4.2)$$

A classical result of Steinberg implies that $\mathcal{X}'_\omega(M)_\lambda$ is finite dimensional. Since D_E^\pm commute with the \mathbb{H} -action, we have the restricted Dirac operators

$$D_E^\pm(\lambda) : \mathcal{X}'_\omega(E \otimes S^\pm)_\lambda \rightarrow \mathcal{X}'_\omega(E \otimes S^\mp)_\lambda, \quad (6.4.3)$$

and the restricted Dirac index

$$I_E(\lambda) = \ker D_E^+(\lambda) - \ker D_E^-(\lambda), \quad (6.4.4)$$

a virtual \mathbb{H} -module. We will see that $I_E(\lambda) = 0$ except for finitely many values of λ , depending on E . We have the following easy restriction.

Lemma 6.11. *Assume $\tilde{\delta}$ is an irreducible \widetilde{W}' -representation. Then $\ker D_{\tilde{\delta}}^{\pm}(\lambda) \neq 0$ only if $\langle \lambda, \lambda \rangle = \langle \chi^{\tilde{\delta}}, \chi^{\tilde{\delta}} \rangle$, where $\chi^{\tilde{\delta}}$ is the central character attached to $\tilde{\delta}$ by Definition 4.5.*

Proof. Assume that $\ker D_{\tilde{\delta}}^+(\lambda) \neq 0$. Then $\ker D_{\tilde{\delta}}^- D_{\tilde{\delta}}^+(\lambda) \neq 0$, and by (6.3.2) and (6.3.8), we get that $\langle \lambda, \lambda \rangle = \tilde{\delta}(\Omega_{\widetilde{W}}) = \langle \chi^{\tilde{\delta}}, \chi^{\tilde{\delta}} \rangle$. \square

In the next section, we show that in fact, under the assumptions of Lemma 6.11, $\lambda = \chi^{\tilde{\delta}}$, as a consequence of Theorem 3.2.

Lemma 6.12. *For every $\lambda \in W \setminus V_{\mathbb{C}}^{\vee}$, $I_E(\lambda) = \mathcal{X}'_{\omega}(E \otimes S^+)_{\lambda} - \mathcal{X}'_{\omega}(E \otimes S^-)_{\lambda}$.*

Proof. We have $(x, D_E(\lambda)^- y)_{k, C_+} = (D_E^+(\lambda)x, y)_{k, C_+} = 0$, whenever $x \in \ker D_E^+(\lambda)$, which shows that $\ker D_E^+(\lambda) \subset (\operatorname{im} D_E^-(\lambda))^{\perp}$. Conversely, if $x \in (\operatorname{im} D_E^-(\lambda))^{\perp} \subset \mathcal{X}'_{\omega}(E \otimes S^+)_{\lambda}$, then $0 = (x, D_E^-(\lambda)y)_{k, C_+} = (D_E^+(\lambda)x, y)_{k, C_+}$, for all $y \in \mathcal{X}'_{\omega}(E \otimes S^-)_{\lambda}$. Specializing to $y = D_E^+(\lambda)x$, we see that $D_E^+(\lambda)x = 0$, hence $x \in \ker D_E^+(\lambda)$. Therefore, we proved

$$\begin{aligned} \mathcal{X}'_{\omega}(E \otimes S^+)_{\lambda} &= \ker D_E^+(\lambda) \oplus \operatorname{im} D_E^-(\lambda), \text{ and similarly,} \\ \mathcal{X}'_{\omega}(E \otimes S^-)_{\lambda} &= \ker D_E^-(\lambda) \oplus \operatorname{im} D_E^+(\lambda). \end{aligned}$$

Then $D_E^+(\lambda)$ maps $\operatorname{im} D_E^-(\lambda)$ onto $\operatorname{im} D_E^+(\lambda)$ and $D_E^-(\lambda)$ maps $\operatorname{im} D_E^+(\lambda)$ onto $\operatorname{im} D_E^-(\lambda)$, and they induce an isomorphism $\operatorname{im} D_E^+(\lambda) \cong \operatorname{im} D_E^-(\lambda)$ as \widetilde{W}' -representations. The claim follows. \square

Definition 6.13. For a given genuine \widetilde{W}' -representation E , we define the global Dirac index to be

$$I_E = \bigoplus_{\lambda} I_E(\lambda), \quad (6.4.5)$$

where $I_E(\lambda)$ is as in (6.4.4). In fact, we conjecture that $\ker D_E^+$ and $\ker D_E^-$ are finite dimensional, and as a result, that $I_E = \ker D_E^+ - \ker D_E^-$.

6.5. A realization of discrete series.

Lemma 6.14. *Let (π, X) be an irreducible \mathbb{H} -module and M a finite dimensional W' -module such that $\operatorname{Hom}_{\mathbb{H}}(X, \mathcal{X}'_{\omega}(M)) \neq 0$. Then (π, X) is a discrete series \mathbb{H} -module.*

Proof. Let $x \in X$ be an $S(V_{\mathbb{C}})$ -weight vector in X with weight $\nu \in V^{\vee}$. This means that for all $\xi \in V_{\mathbb{C}}$, $\pi(\xi)x = (\xi, \nu)x$. Let $\kappa \in \operatorname{Hom}_{\mathbb{H}}(X, \mathcal{X}'_{\omega}(M))$ be nonzero. Then

$$\partial(\xi)\kappa(x) = \kappa(\pi(\xi)x) = (\xi, \nu)x.$$

Recall that in $\mathcal{X}'_{\omega}(M)$, $\partial(\xi)$ acts by differentiation only on the function part of the tensor product, so we may assume without loss of generality, that $\kappa(x) = f \otimes m$, for some $f \in C^{\omega}(V)$, $m \in M$. This means that the one-dimensional vector space

generated by f is invariant under differentiation by all elements $p \in S(V_{\mathbb{C}})$, and therefore f is an exponential function $f(v) = ce^{(v, \nu)}$, $v \in V$.

But since $f \otimes m \in \mathcal{X}'_{\omega}(M)$, f must be an L^2 -function on C_+ , and therefore the weight ν must satisfy the strict Casselman criterion (see 2.3.2). \square

Lemma 6.15. *Suppose that (π, X) is an irreducible discrete series \mathbb{H} -module and M a finite dimensional W' -module. Then*

$$\mathrm{Hom}_{\mathbb{H}}(X, \mathcal{X}'_{\omega}(M)) \cong \mathrm{Hom}_{W'}(M^*, X^*). \quad (6.5.1)$$

Proof. Since the dual of $S(V_{\mathbb{C}})$ can be identified with Laurent series on $V_{\mathbb{C}}$, the proof of Lemma 6.14 shows that every nontrivial homomorphism of X into $(\mathbb{H} \otimes_{W'} M^*)^*$, under the assumption that X be a discrete series, lands in fact in $\mathcal{X}'_{\omega}(M)$. In other words, we have

$$\mathrm{Hom}_{\mathbb{H}}(X, \mathcal{X}'_{\omega}(M)) \cong \mathrm{Hom}_{\mathbb{H}}(X, (\mathbb{H} \otimes_{W'} M^*)^*).$$

Furthermore, using the tautological isomorphism $\mathrm{Hom}(A, B^*) \cong \mathrm{Hom}(B, A^*)$, we find that

$$\begin{aligned} \mathrm{Hom}_{\mathbb{H}}(X, \mathcal{X}'_{\omega}(M)) &\cong \mathrm{Hom}_{\mathbb{H}}(\mathbb{H} \otimes_{W'} M^*, X^*) \\ &\cong \mathrm{Hom}_{W'}(M^*, X^*), \end{aligned} \quad (6.5.2)$$

where the last step is the Frobenius isomorphism. \square

Recall that by Corollary 2.10, if (π, X) is a discrete series, then $r(X)$ is a unit element in $\overline{R}_{\mathbb{Z}}(W)$, where r is the restriction map from 2.6.1. Therefore, by Theorem 4.2(2), there exists unique irreducible \widetilde{W}' -representations $\widetilde{\delta}_X^+, \widetilde{\delta}_X^-$ such that

$$I(X) = \widetilde{\delta}_X^+ - \widetilde{\delta}_X^-, \quad (6.5.3)$$

and moreover, by Corollary 4.6, the central character of X is $\chi^{\widetilde{\delta}_X^+} = \chi^{\widetilde{\delta}_X^-}$, where the notation is as in Definition 4.5.

Theorem 6.16. *For any irreducible genuine \widetilde{W}' -module E , I_E (as defined by (6.4.5)) is a virtual finite linear combination of discrete series representations. Let (π, X) be an irreducible discrete series with central character $\chi^{\widetilde{\delta}_X^+}$ and let E be an irreducible \widetilde{W}' -representation. Then*

$$\mathrm{Hom}_{\mathbb{H}}(X, I_E) \cong \mathrm{Hom}_{\mathbb{H}}(X, I_E(\Lambda(\widetilde{\delta}_X))) \cong \mathrm{Hom}_{\widetilde{W}'}(E^*, I(X)), \quad (6.5.4)$$

and the dimension of these spaces is $\begin{cases} 1, & \text{if } E^* \cong \widetilde{\delta}_X^+ \\ -1, & \text{if } E^* \cong \widetilde{\delta}_X^- . \text{ In particular, } I_{\widetilde{\delta}_X^+} \text{ is the} \\ 0, & \text{otherwise.} \end{cases}$

unique irreducible discrete series \mathbb{H} -module X that has Dirac index $\widetilde{\delta}_X^+ - \widetilde{\delta}_X^-$.

Proof. Since the central character of X is $\Lambda(\widetilde{\delta}_X)$, we necessarily have $\mathrm{Hom}_{\mathbb{H}}(X, I_E) \cong \mathrm{Hom}_{\mathbb{H}}(X, I_E(\Lambda(\widetilde{\delta}_X)))$. By Lemma 6.12, this space is isomorphic with

$$\mathrm{Hom}_{\mathbb{H}}(X, \mathcal{X}'_{\omega}(E \otimes S^+) - \mathcal{X}'_{\omega}(E \otimes S^-)),$$

which by Lemma 6.15 equals

$$\mathrm{Hom}_{W'}(E^* \otimes (S^+ - S^-)^*, X^*) = \mathrm{Hom}_{\widetilde{W}'}(E^*, X \otimes (S^+ - S^-)) = \mathrm{Hom}_{\widetilde{W}'}(E^*, I(X));$$

here, we used that $X^* \cong X$ as W' -modules, while the last equality is from Lemma 4.1. Now (6.5.4) follows from this using that $I(X) = \widetilde{\delta}_X^+ - \widetilde{\delta}_X^-$.

To complete the proof, notice that Theorem 4.2(2) combined with Corollary 2.10 imply that if $X \not\cong Y$ are two distinct irreducible discrete series, then $\tilde{\delta}_X^\pm \neq \tilde{\delta}_Y^\pm$, and therefore X is the only discrete series that contributes to $I_{\tilde{\delta}_X^\pm}^\pm$. \square

Remark 6.17. If instead of the inner product $(\ , \)_{k, C_+}$, one uses $(\ , \)_{k, C_-}$ (see Remark 6.4), then one can realize all irreducible anti-discrete series of \mathbb{H} (i.e., the Iwahori-Matsumoto duals of the discrete series) in the corresponding analytic models and indices of Dirac operators. All the results are the obvious analogues.

Remark 6.18. Most of the constructions here apply equally well to noncrystallographic root systems. Theorem 6.16 shows that if X is an irreducible discrete module such that $X|_W$ is a unit element in $\overline{R}_Z(W)$ with respect to the elliptic pairing, then X is isomorphic to the global Dirac index for an irreducible \widetilde{W} -representation, and in particular, it is unitary.

REFERENCES

- [AS] M. Atiyah, W. Schmid, *A geometric construction of the discrete series for semisimple Lie groups*, Invent. Math. **42** (1977), 1–62.
- [BCT] D. Barbasch, D. Ciubotaru, P. Trapa, *The Dirac operator for graded affine Hecke algebras*, to appear in Acta Math.
- [BM] D. Barbasch, A. Moy, *A unitarity criterion for p-adic groups*, Invent. Math. **98** (1989), 19–38.
- [BS] W. Beynon, N. Spaltenstein, *Green functions of finite Chevalley groups of type E_n ($n = 6, 7, 8$)*, J. Algebra **88** (1984), no. 2, 584–614.
- [CT] D. Ciubotaru, P. Trapa, *Characters of Springer representations on elliptic conjugacy classes*, preprint, [arXiv:1105.4113v2](https://arxiv.org/abs/1105.4113v2).
- [Ca] R.W. Carter, *Finite groups of Lie type*, Pure and Applied Math. (New York), Wiley-Interscience, NY (1985), xii+544pp.
- [C] D. Ciubotaru, *Spin representations of Weyl groups and Springer’s correspondence*, J. Reine Angew. Math., doi: 10.1515/crelle.2011.160.
- [CK] D. Ciubotaru, S. Kato, *Tempered modules in the exotic Deligne-Langlands classification*, Adv. Math. **226** (2011), 1538–1590.
- [CKK] D. Ciubotaru, M. Kato, S. Kato, *On characters and formal degrees of discrete series of classical affine Hecke algebras*, Invent. Math., doi: 10.1007/s00222-011-0338-3.
- [DO] P. Delorme, E. Opdam, *Schwartz algebra of an affine Hecke algebra*, J. Reine Angew. Math. **625** (2008), 59–114.
- [HO] G.J. Heckman, E.M. Opdam, *Yang’s system of particles and Hecke algebras*, Ann. Math. **45** (1997), 139–173.
- [EOS] E. Emsiz, E.M. Opdam, J.V. Stokman, *Trigonometric Cherednik algebra at critical level and quantum many-body problems*, Selecta Math. (N.S.) **14** (2009), no. 3-4, 571–605.
- [K] S. Kato, *An exotic Deligne-Langlands correspondence for symplectic groups*, Duke Math. J. **148**(2), 305–371 (2009).
- [KL] D. Kazhdan, G. Lusztig, *Proof of Deligne-Langlands conjecture for Hecke algebras*, Invent. Math. **87** (1987), 153–215.
- [La] V. Lafforgue *Banach KK-theory and the Baum-Connes Conjecture*, ICM 2002 Vol. III **13**, 795–811.
- [L1] G. Lusztig, *Affine Hecke algebras and their graded version*, J. Amer. Math. Soc. **2** (1989), no. 3, 599–635.
- [L2] G. Lusztig, *Cuspidal local systems and graded Hecke algebras III*, Represent. Theory **6** (2002), 202242.
- [M] A. Morris, *Projective characters of exceptional Weyl groups*, J. Algebra **29** (1974), 567–586.
- [O1] E.M. Opdam, *On the spectral decomposition of affine Hecke algebras*, J. Math. Jussieu **3** (2004), no. 4, 531–648.

- [O2] E.M. Opdam, *The central support of the Plancherel measure of an affine Hecke algebra*, Mosc. Math. J. 7 (2007), 723–741, 767–768.
- [R] M. Reeder, *Euler-Poincaré pairings and elliptic representations of Weyl groups and p -adic groups*, Compositio Math. 129 (2001), no. 2, 149–181.
- [Re] E.W. Read, *On projective representations of the finite reflection groups of type B_l and D_l* , J. London Math. Soc. (2), 10 (1975), 129–142.
- [OS1] E. Opdam, M. Solleveld, *Homological algebra for affine Hecke algebras*, Adv. Math. 220 (2009), no. 5, 1549–1601.
- [OS2] E. Opdam, M. Solleveld, *Discrete series characters for affine Hecke algebras and their formal degrees*, Acta Math. 205 (2010), 105–187.
- [Sl] K. Slooten, *Induced discrete series representations for Hecke algebras of types B_n^{aff} and C_n^{aff}* , Int. Math. Res. Not. IMRN 2008, no. 10, Art. ID rnn023, 41 pp.
- [So] M. Solleveld, *On the classification of irreducible representations of affine Hecke algebras with unequal parameters*, Represent. Theory 16 (2012), 1–87.

(D. Ciubotaru) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF UTAH, SALT LAKE CITY, UT 84112, USA

E-mail address: ciubo@math.utah.edu

(E. Opdam) KORTEWEG-DE VRIES INSTITUTE FOR MATHEMATICS, UNIVERSITEIT VAN AMSTERDAM, SCIENCE PARK 904, 1098 XH AMSTERDAM, THE NETHERLANDS

E-mail address: e.m.opdam@uva.nl

(P. Trapa) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF UTAH, SALT LAKE CITY, UT 84112, USA

E-mail address: ptrapa@math.utah.edu